

AD-A147 281

CONFERENCE ON SINGLET MOLECULAR OXYGEN (COSMO 84) HELD  
AT CLEARWATER BEAC. (U) UNIVERSITY OF SOUTH FLORIDA  
TAMPA B STEVENS 24 SEP 84 18-2705-124F


1/1


UNCLASSIFIED

N00014-84-C-0032

F/G 7/2


NL





AD-A147 281

# CONFERENCE ON SINGLET MOLECULAR OXYGEN



JANUARY 4-7, 19**84**  
CLEARWATER, BEACH, FLORIDA

PROGRAM  
and  
ABSTRACTS

DTIC FILE COPY

A  
N00014-84C-0032

This document has been approved  
for public release and sale; its  
distribution is unlimited.

84 : 10 01 041

FINAL REPORT

on

CONFERENCE ON SINGLET MOLECULAR OXYGEN (COSMO 84)  
(Report #18-2705-124F)

by

BRIAN STEVENS

UNIVERSITY OF SOUTH FLORIDA, TAMPA, FL. 33620

to

OFFICE OF NAVAL RESEARCH CODE:612B:PMW  
800 NORTH QUINCY STREET  
ARLINGTON, VA 22217

Security Classification - Unclassified

Number of Pages - 44

Distribution: -

Associate Director for Life Science Programs ONR (1)  
ACO, ONR, Georgia Institute of Technology (1)  
Defense Technical Information Center (1)

Award Amount - \$5000

Duration - 1-1-84 to 6-30-84

Contract # N00014-84-C-0032

Signature



Brian Stevens

Date:

September 24, 1984

CONFERENCE ON SINGLET MOLECULAR OXYGEN  
COSMO 84  
January 3-7th, 1984  
Clearwater Beach, Florida

Some 80 scientists from 12 countries attended the Conference at which 47 papers were presented in the following sessions:

Sources and Yields (B. Stevens)  
Relaxation and Quenching (M. A. J. Rodgers)  
Reaction Mechanisms (C. S. Foote)  
Biological/Biomedical Implications (R. W. Murray)  
Gas Phase and Atmospheric Manifestations (R. P. Wayne)  
Synthetic and Other Applications (A. P. Schaap)

The sessions organizers indicated, together with A. M. Trozzolo, formed the planning committee.

Sponsors included:

The Office of Naval Research  
The National Science Foundation  
The American Society for Photobiology  
The Inter-American Photochemical Society  
Jacobs-Fendig, Inc.  
Mr. Robert Dressler  
The University of South Florida

Microns

Superscript 1

Superscript 1

Delta g

The proceedings will be published in a dedicated issue (3 volumes) of the Journal of Photochemistry later this year.

The Conference objective was to promote interdisciplinary awareness and communication by assembling research workers in such diverse fields as photochemistry, photophysics, synthetic chemistry, photobiology, photomedicine, laser and atmospheric physics, to present their recent findings with singlet molecular oxygen as the common theme. This was prompted by several recent developments, notably the direct observation of  $O_2^1\Delta_g$  in emission at 1.27  $\mu$  which now provides solvent-dependent lifetimes, reaction or quenching rate constants, and sensitized yields directly, the trapping of zwitterionic intermediates and the role of catalysts in electron transfer peroxidation, the feasibility of a singlet oxygen-iodine chemical laser, direct observation of  $O_2^1\Delta_g$  in enzymic processes, the use of endoperoxides as actinometers, the phototherapy of malignant tumors, and prospects for  $O_2^1\Delta_g$  as a solar energy storage intermediate. If attendance on the final morning (which equalled that at the opening session) is a criterion of success, then any failure to meet the stated objectives was not the fault of the participants. Since COSMO 84 does not represent the continuation of a Conference series, e.g. on Oxygen radicals (Pinawa 1977, Texas 1980, Munich 1983), on Polymers (Stockholm 1976) or the role of singlet oxygen in environmental sciences (New York 1970) more objective criteria of its success will be provided by the reception of the Conference Proceedings and the planning of COSMO 87.

SPONSORSHIP  
OF THE  
CONFERENCE ON SINGLET MOLECULAR OXYGEN  
COSMO 84  
by

THE OFFICE OF NAVAL RESEARCH  
(Division of Biological Sciences)  
THE NATIONAL SCIENCE FOUNDATION  
(Biophysics and Atmospheric Science Programs)  
THE INTERAMERICAN PHOTOCHEMICAL SOCIETY  
THE AMERICAN SOCIETY FOR PHOTOBIOLOGY  
JACOBS-FENDIG INC.  
MR. ROBERT DRESSLER  
THE UNIVERSITY OF SOUTH FLORIDA

IS GRATEFULLY ACKNOWLEDGED

---

ORGANIZING COMMITTEE

BRIAN STEVENS (HOST AND CO-ORDINATOR)  
CHRISTOPHER S. FOOTE  
ROBERT W. MURRAY  
MICHAEL A.J. RODGERS  
A. PAUL SCHAAP  
ANTHONY M. TROZZOLO  
RICHARD P. WAYNE

---

1V00014-84C-0032

These Abstracts are not to be considered as publications  
of the work described and should not be cited.

CONFERENCE AND PROGRAM NOTES

1. ALL SESSIONS will be HELD in the BIG SURF BALL ROOM.
2. REGISTRATION will take place at the RECEPTION on TUESDAY EVENING and in the HOTEL LOBBY throughout the CONFERENCE.
3. SPEAKERS are asked to give their SLIDES to one of the PROJECTIONISTS before the appropriate session and their MANUSCRIPTS to the REGISTRATION DESK before they leave.
4. In order to avoid parallel and poster sessions it has been necessary to schedule MORNING, AFTERNOON and EVENING SESSIONS on WEDNESDAY.
5. Since the attendance of some PROGRAMMED SPEAKERS is UNCERTAIN at the time of 'going to press', there is NO TIMETABLE. COFFEE, LUNCH and (on WEDNESDAY) DINNER BREAKS will be announced by the SESSION CHAIRMAN.
6. LOCAL INFORMATION may be obtained at the HILTON REGISTRATION DESK or the LOBBY REGISTRATION DESK. A LINDO'S TOUR DESK is located in the HILTON LOBBY and a LINDO'S AGENT has been invited to the RECEPTION on TUESDAY.
7. Please note that in order to BOARD the COACHES on THURSDAY EVENING at 6:30 PM you will need a TICKET for the LUAU DINNER.
8. You are reminded that the WEATHER can be COOL and WET on the BEACH in JANUARY.



*Auto enfil*

*A-1*

# PROGRAM

TUESDAY EVENING

JANUARY 3

7 - 10 PM

## RECEPTION AND REGISTRATION

Poolside (weather permitting) or Crown Ballroom

WEDNESDAY MORNING/AFTERNOON JANUARY 4

Big Surf Ballroom

8:25 AM

WELCOMING REMARKS AND ANNOUNCEMENTS

8:30 AM

## SESSION on SOURCES and YIELDS

B. STEVENS - Chairman

- |                 |    |   |
|-----------------|----|---|
| A. SINGH        | S5 | 'Possible Formation of Singlet Oxygen from Vibrationally-Excited Water'                               |
| J.R. KANOFKY    | K1 | 'Singlet Oxygen Production by Lactoperoxidase: Halide Dependence and Quantitation of Yield'           |
| D.F. EVANS      | E2 | 'Studies on Singlet Oxygen in Aqueous Solution'   |
| G.P. GURINOVICH | G6 | 'Mechanism of Singlet Oxygen Luminescence Sensitization by Polyatomic Molecules'                      |
| A.A. GORMAN     | G2 | 'The Efficiency of Triplet Energy Transfer to $O_2(^3\Sigma_g^-)$ : Sensitizer and Medium Dependence' |
| C. TANIELIAN    | T1 | 'Production and Quenching of Singlet Oxygen by the Sensitizer in Dye-Sensitized Photooxygenations'    |

## SESSION on RELAXATION and QUENCHING

M.A.J. Rodgers and P.R. Ogilby - Chairmen

- |                 |    |  |
|-----------------|----|--|
| M.A.J. RODGERS  | R2 | 'The Relaxation of $O_2(^1\Delta_g)$ in Condensed Systems'   |
| M. MAIER        | M1 | 'Relaxation of the $^1\Delta_g$ State in Pure Liquid Oxygen and in Liquid Mixtures of $^{16}O_2$ and $^{18}O_2$ '                      |
| J.G. PARKER     | P1 | 'Optical Determination of the Rates of Formation and Decay of $O_2(^1\Delta_g)$ in $H_2O$ , $D_2O$ and Other Solvents'                 |
| B. STEVENS      | S7 | 'Preferential Solvation of $O_2(^1\Delta_g)$ in Mixed Solvents'  |
| B.M. DZHAGAROV  | D2 | 'Singlet Oxygen Luminescence'  |
| A.A. KRASNOVSKY | K3 | 'Photosensitized Luminescence of Singlet Oxygen in Solutions: Mechanism of Quenching by Solvents and Biologically Important Compounds' |
| F. WILKINSON    | W5 | 'The Mechanism of Quenching of Singlet Oxygen by Molecular Iodine'   |



WEDNESDAY AFTERNOON/EVENING  
SESSION ON REACTION MECHANISMS  
C.S. FOOTE - Chairman

W. ANDO	W1	'Stereochemical Studies on the Mechanism of Dioxetane and Epoxide Formation from Hindered Olefins'
A.P. SCHAAP	S3	'Nucleophilic Oxygen - Atomic Transfer from a Perepoxy Intermediate to Sulfoxides'
G.W. GRIFFIN	G3	'Sensitized Photooxidation of Methyl Substituted Cyclobutenes'
A.P. SCHAAP	S2	'Cosensitized Electron - Transfer Photo-oxygenation'
P.B. GANDHI	G1	'Dye-sensitized Photo-oxygenation of Isoniazid by Singlet Molecular Oxygen'
J.S. CONNOLLY	C1	'Effects of Solvent on the Rate of Bacterio-chlorophyll <u>a</u> Photooxidation'
I. FORSSKAHL	F3	'Aspects of Photosensitized Lignin Oxygenation'
A.A. FRIMER	F4	'The effect of Strain on the Rearrangements of Primary Singlet Oxygen Products'
G.D. MENDENHALL	M2	'Structure and Kinetics of the Uncatalyzed Decomposition of Trialkyl Phosphite Ozonides'

THURSDAY

JANUARY 5

SESSION ON BIOLOGICAL/BIOMEDICAL IMPLICATIONS

8:30 AM

R.W. MURRAY - Chairman

- D.G. WHITTEN W4 'Self-Sensitized Photooxidation of Protoporphyrin IX and Related Porphyrins in Erythrocyte Ghosts and Microemulsions: A Novel Photooxidation Pathway'
- L.I. GROSSWEINER G4 'Reactions of Singlet Oxygen with Liposomes'
- R.B. CUNDALL C2 'Oxidation Photosensitized by 2-Chlorothioxanthone: The Role of Singlet Oxygen'

BREAK

L.I. GROSSWEINER - Chairman

- M.K. LOGANI L2 't-Butylated Hydroxytoluene Sensitized Photooxidation of the Fatty Acids: Inhibition by Diazabicyclooctane and Role of Singlet Oxygen'
- N. DURAN D1 'Singlet Oxygen Generation from the Peroxidase-Catalysed Aerobic Oxidation of an Activated - CH<sub>2</sub> - Substrate'
- C.S. FOOTE F2 'Singlet Oxygen Production from Photodynamic Sensitizers'

LUNCH

L.L. SMITH - Chairman

- H. SINGH S6 'Singlet Oxygen and Ribosomes: Inactivation and Sites of Damage'
- M.J. PEAK P3 'Reactive Oxygen Species in the Biological Effects of Solar Ultraviolet Radiation'
- S. ZIGMAN Z1 'Active Oxygen Chemistry of Ocular Tissues'

BREAK

A.M. TROZZOLO - Chairman

- W. NULTSCH N1 'Possible Role of Singlet Oxygen in the Control of the Phototactic Reaction Sign of the Cyanobacterium Anabaena Vaniabilis'
- A.U. KHAN K2 ' $O_2(^1\Delta_g)$  Emission from Enzyme Reactions'
- R.W. MURRAY M3 'Singlet Oxygen Oxidation of Substituted Thiobenzamides'

6:30 PM

COACHES LEAVE HILTON FOR TIKI GARDENS

FRIDAY

JANUARY 6

8:30 AM

SESSION ON GAS PHASE AND ATMOSPHERIC MANIFESTATIONS

R.P. WAYNE - Chairman

R.P. WAYNE	W2	'Laboratory Kinetics and Spectroscopy Applied to Airglow Studies'
E.J. LLEWELLYN	L1	'Singlet Molecular Oxygen in Planetary Atmospheres'
E.A. OGRYZLO	O1	'The Yield of $O_2(b^1\Sigma_g^+)$ in Oxygen Atom Re-combination'
P. BORRELL	B2	'High Temperature Kinetic and Spectroscopic Studies of Singlet Oxygen'
J.G. PARKER	P2	'Optical Determination of $O_2(^1\Delta_g)$ Quenching Rates and Relative Emission Intensities in High Pressure Oxygen Gas Using Pulsed Laser Excitation at 10640 Å'
D.J. BOGAN	B1	'Chemiluminescent Reactions of Organic Molecules Forming or Consuming Singlet Oxygen'
E.H. FINK	F1	' $O_2(^1\Delta_g)$ Sensitized Chemiluminescence of $a^1\Delta - X^3\Sigma^-$ and $b^1\Sigma^+ - X^3\Sigma^-$ Transitions of Group V-VII Diatomic Molecules'
W.C. EISENBERG	E1	'Generation and Reactions of Singlet Delta Oxygen at Ambient Atmospheric Conditions'
R.F. HEIDNER, III	H1	'Behaviour of Singlet Oxygen in the Oxygen-Iodine Transfer Laser'
P.L. HOUSTON	H2	'Kinetics of the Chemical Oxygen-Iodine Laser'
P.D. WHITEFIELD	W3	' $O_2(^1\Delta_g)$ as an Energy Transfer Partner for Chemical Laser Pumping ( $O_2^1\Delta + IF$ )'

SATURDAY MORNING

JANUARY 7

8:30 AM

SESSION ON SYNTHETIC AND OTHER APPLICATIONS

A.P. SCHAAP - Chairman

- |                |    |   |
|----------------|----|---|
| H.H. WASSERMAN | W1 | 'Recent Applications of Singlet Oxygen Reactions in Synthesis'  |
| H.-D. BRAUER   | B3 | 'Photoperoxidation of Aromatic Compounds and Wavelength - Dependent Photochemistry of the Obtained Peroxides'                         |
| R. SCHMIDT     | S4 | 'Self-Sensitized Photooxidation of Aromatic Compounds and Photocycloreversion of Endoperoxides. Applications in Chemical Actinometry' |
| J.E. GUILLET   | G5 | 'Prospects for Solar Synthesis. I. A New Method for Singlet Oxygen Reactions Using Natural Sunlight'                                  |
| I. SAITO       | S1 | 'On the Use of Water Soluble $^1\text{O}_2$ Sources for Mechanistic Study and for Oxygenation of Biological Systems'                  |
| J.F. RABEK     | R1 | 'Synergistic and Antagonistic Effects of Quenching Singlet Oxygen in a Polymer Matrix'  |
| C.W. JEFFORD   | J1 | 'Construction of 1,2,4-Trioxane Rings by Photooxygenation of Enol Ethers'   |

CLOSING REMARKS

Stereochemical Studies on the Mechanism of  
Dioxetane and Epoxide Formations from Hindered Olefins

Wataru Ando, Yoshio Kabe and Toshikazu Takata

Department of Chemistry, University of Tsukuba


Sakura-mura, Ibaraki 305, Japan

Katsuhiko Ueno



Research Institute for Polymers and Textile

Yatabe-cho, Ibaraki 305, Japan


Considerable interest is focused on reaction of sterically hindered olefins such as adamantylidene adamantane (1), since oxygenation of 1 gives the remarkably stable dioxetane sometimes with accompanying formation of epoxides. Such observation has



enlivened the controversy that deoxygenation of peroxide occurs or not. More recently, it has been shown that cation radical of 1 uniquely reacts with triplet oxygen to yield dioxetane by chain reaction. In spite of the continuous interests concerning with hindered olefins, no intrinsic stereochemical study has been done yet. Then, we synthesized two stereoisomeric di-tert-butyl (bicyclo[3,3,1]non-9-ylidene)s (2-anti and 2-syn). Assignments of the stereochemistry followed from the X-ray crystal analysis of 2-anti (Fig. 1). The first objective is on the mechanism of dioxetane formation. Either TPP (tetraphenylporphine) or DCA




(9,10-dicyanoanthracene)-sensitized photooxygenation of 2-anti gave stereospecifically single dioxetane 3-cis,trans. Similarly, photooxygenation of 2-syn by either TPP or DCA gave two dioxetanes, 3-cis,cis and 3-trans,trans, respectively. Structures of all dioxetanes (3) isolated were easily assigned by spectroscopies



and X-ray crystal analysis of 3-cis,cis (Fig. 2). In contrast, the electrode-catalysed oxygenation of each isomeric olefin gave mixture of all three dioxetanes. These results suggested that the concerted cycloaddition of singlet oxygen occurred in dye-sensitized oxygenation, and singlet oxygen might also be involved in the case of DCA. In the electrode-catalysed oxygenation, cation radical is seriously anticipated.

Second set of experiments is concerned with photoepoxidation. Both Rose bengal and TPP-sensitized photooxygenations of 2 in the presence of methyl phenyl sulfoxide gave rise to the formation of epoxides (4) with retention of stereochemistry. These results



might confirm the deoxygenation of peroxide. Meanwhile, biacetyl-sensitized photoepoxidation has been known to be non-stereospecific. However, the photoepoxidation of 2 was completely stereospecific. Peracid formed by hydrogen abstraction of acylperoxy radical may be actual oxidizing species.

B1  
Friday

Chemiluminescent Reactions of Organic  
Molecules Forming or Consuming  
Singlet Oxygen

Denis J. Bogan  
Combustion & Fuels Branch  
Code 6180  
Naval Research Laboratory  
Washington, D. C. 20375

This paper will present and discuss gas phase chemiluminescence studies of two reactions, each proceeding through a long lived vibrationally excited peroxide intermediate. These are;

- (1) Reaction of  $O_2 (^1\Delta_g)$  with olefin giving electronically excited carbonyl product via a dioxetane intermediate, and
- (2) Reaction of two (non-tertiary) alkyl peroxy radicals by Russell's mechanism giving  $O_2 (^1\Sigma_g)$  via a dialkyl tetraoxide intermediate. Reaction coordinate diagrams for typical examples of (1) and (2) are shown below.




Figure 1. Schematic Reaction Coordinate Diagram for Reaction (1)




Figure 2. Schematic Reaction Coordinate Diagram for Reaction (2) Enthalpy scale is for the isopropyl peroxy system.

We have studied many examples of reaction (1) in a flow system where  $O_2$  ( ${}^1\Delta$ ) was formed by 2450 MHz discharge of  $O_2$  doped with a trace of Hg vapor to recombine O atoms. Extensive data on product energy partitioning have been obtained from chemiluminescence spectra.

Examples of reaction (2) have also been studied in the flow system. Trapping and analysis were performed for all products of reacting mixtures typically consisting of; 98%  $O_2$ , less than 1/2% O ( ${}^3P$ ) and/or Cl ( ${}^2P$ ) atoms, and 1 to 1.5% RH, at temperatures (200 - 450  $^{\circ}C$ ) chosen to insure the loss of alkoxy radicals by unimolecular fragmentation. The presence of  $O_2$  ( ${}^1\Sigma$ ) product was confirmed by the spectrum of  $O_2$  ( ${}^1\Sigma \rightarrow {}^3\Sigma$ ) chemiluminescence in the isopropyl peroxy reaction. This experiment made  $iPrO_2^{\cdot}$  via reaction of H with dimethyl ketene in the presence of a large excess of  $O_2$ . The combined results of our studies of reaction (2) are interpreted as proof of the occurrence of Russell's mechanism in the gas phase.



For presentation at COSMO '84

High Temperature Studies of Singlet Molecular Oxygen

BY

Peter Borrell, Patricia M. Borrell, D.S. Richards and Razmik Boodaghians  
Department of Chemistry, Keele University, Staffordshire, England.

The aim of the work to be presented is to study the collisional quenching reactions of  $O_2(a^1\Delta_g)$  and  $O_2(b^1\Sigma_g^+)$  over as wide a temperature range as possible with a variety of quenching partners. The paper reviews the experimental technique and gives an account of the principal results obtained for both species including some new results for quenching by hydrogen.

The method used is a combined discharge flow/shock tube technique in which singlet oxygen is generated in a microwave discharge and then heated in a glass shock tube. The factors which contribute to the reliability of the method for obtaining results between 600 and 1800 K will be discussed.

For the quenching of  $O_2(a^1\Delta_g)$  there are two classes of quencher: those for which there is an Arrhenius dependence of rate constant on temperature with an activation energy between 15 and 20  $\text{kJ mol}^{-1}$ , HCl, NO and  $\text{SO}_2$ , and those for which the change is not measurable in our system ( $E < 8 \text{ kJ mol}^{-1}$ ).

There are again two classes of quencher for  $O_2(b^1\Sigma_g^+)$ ; poor quenchers at room temperature,  $\text{N}_2$ ,  $\text{O}_2$ ,  $\text{SO}_2$  for which the rate constant increases with temperature; and more efficient quenchers, HCl, NO,  $\text{NH}_3$ ,  $\text{CO}_2$ ,  $\text{N}_2\text{O}$  for which the rate constant is nearly independent of temperature or actually decreases above 1000 K.

The present theoretical approaches do not appear to be capable of explaining these results in a quantitative way.

PHOTOPEROXIDATION OF AROMATIC COMPOUNDS AND WAVELENGTH-  
DEPENDENT PHOTOCHEMISTRY OF THE OBTAINED ENDOPEROXIDES

H.-D. Brauer, A. Acs, W. Drews, R. Gabriel, S. Ghaeni  
K. Schaffner, W. Trost and R. Schmidt

Institute for Physical and Theoretical Chemistry,  
University of Frankfurt, Robert-Mayer-Str. 11,  
D 6000 Frankfurt/Main, F.R.G.

A. Photoperoxidation of Aromatic Compounds

A.1. Generation of  $^1\text{O}_2$

It is well known that  $^1\text{O}_2$  in its  $^1\Delta_g$ -state is the reactive intermediate in the sensitized and self-sensitized photoperoxidation of aromatic compounds leading to endoperoxide formation.  $^1\text{O}_2$  can be generated by both the lowest excited singlet and triplet states of the aromatic compounds, if the singlet-triplet energy separation  $E_{ST}$  and the triplet energies  $E_T$ , respectively, exceed the excitation energy of  $\text{O}_2$   $^1\Delta_g$  at  $7990\text{ cm}^{-1}$ . This was established for the first time for the case of rubrene by measurements of the photoperoxidation quantum yield as a function of the rubrene concentration and of the dissolved oxygen concentration in six solvents [1].

Recently, we have measured the photoperoxidation of a series of

aromatic compounds and evaluated our results in the way described in [1]. From these measurements one can classify the investigated compounds into two groups with respect to oxygen quenching of the  $S_1$ -state.

- I. Oxygen quenching of the  $S_1$ -state of the compounds results exclusively in the formation of the lowest triplet state.
- II. Oxygen quenching of the  $S_1$ -state of the compounds may lead either to the formation of the lowest triplet state or to the formation of the singlet ground state.

Within both groups one can further distinguish two types of compounds with respect to  $^1O_2$  generation. Thus the following subdivision is useful:

- I.1. Each compound molecule in the  $S_1$ -state which is quenched by oxygen generates two molecules of  $^1O_2$ . This is the case for rubrene [1] heterocoerdianthrone [2], 9.10-dimethylantracene [3] and benzodixanthene [3].
- I.2. Each compound molecule in  $S_1$ -state which is quenched by oxygen generates only one molecule of  $^1O_2$ . This is the case for tetracene [3] and anthradichromene [3].
- II.1. Each compound molecule in the  $S_1$ -state which, upon oxy-

gen quenching, passes to the  $T_1$ -state generates two molecules of  $^1O_2$ . This is the case for 9.10-diphenylanthracene [4] and 1.4-dimethoxy-9.10-diphenylanthracene [3]

II.2. Each compound molecule in the  $S_1$ -state which, upon oxygen quenching, passes to the triplet manifold generates only one molecule of  $^1O_2$ . This is the case for meso-diphenylhelianthrene [3].

#### A.2. Wavelength-Dependent Photoperoxidation of Meso-diphenylhelianthrene [3]

Generally, the self-sensitized photoperoxidation of aromatic compounds is independent of the excitation wavelength. As a rule the formation rate of the endoperoxide FR decreases with increasing reaction time, since at a given oxygen concentration the substrate concentration decreases. Meso-diphenylhelianthrene is an exception from this rule.

Excitation with light in the wavelength range between 475 - 610 nm results in behavior similar to that of other aromatic compounds. However, during irradiation at  $\lambda_{exc} \leq 450$  nm FR initially increases remarkably with time reaching a maximum and decreases then continuously. This behavior can be explained taking into account that the formed endoperoxide absorbs stronger than meso-diphenylhelianthrene at the excitation wavelength

and is also able to produce  $^1O_2$ .

B. Wavelength-Dependent Photochemistry of Endoperoxides of Aromatic Compounds

On the basis of orbital and state correlations diagrams, Kearns and Khan predicted a novel type of photoreaction for the photolysis of the endoperoxides of cyclopentadiene and of aromatic compounds, respectively: an adiabatic cycloreversion originating from an upper singlet state. Up to now such a type of photoreaction has never been observed in solution.

The results of our investigations on the photolysis of the endoperoxide of heterococordianthrone confirm for the first time the predictions made by Kearns and Khan [5].

Furthermore, systematic investigations of other endoperoxides lead to the conclusion that the adiabatic cycloreversion is a general phenomenon of the photochemistry of endoperoxides [3,6,7].

Two different reactions are observed during the photolysis of the endoperoxides:

1. Excitation of the  $S_1$ -band causes an irreversible decomposition of the endoperoxides.
2. Excitation of the  $S_2$ -band leads to an adiabatic photocleavage of the endoperoxides into hydrocarbon and  $^1O_2$ .

Some of the systems: endoperoxide  $\xrightleftharpoons[h\nu']{h\nu, \Delta}$  hydrocarbon + O<sub>2</sub> are highly reversible photochromic systems, which are of interest for practical applications [8,9]

# Literature

- [1] H.-D. Brauer and H. Wagener, Ber. Bunsenges. Physik. Chem., 79, 598 (1975)
- [2] H. Wagener and H.-D. Brauer, Mol. Photochem., 7, 441 (1976)
- [3] unpublished data
- [4] W. Drews, R. Schmidt and H.-D. Brauer, Chem. Phys. Lett., 100, 466 (1983).
- [5] R. Schmidt, W. Drews and H.-D. Brauer, J. Am. Chem. Soc., 102, 2971 (1980)
- [6] W. Drews, R. Schmidt and H.-D. Brauer, Chem. Phys. Lett., 70, 84 (1980)
- [7] R. Schmidt, W. Drews and H.-D. Brauer, Z. f. Naturforsch., 37a, 155 (1982)
- [8] H.-D. Brauer, W. Drews and R. Schmidt, J. Photochem., 12, 293 (1980)
- [9] H.-D. Brauer and R. Schmidt, Photochem. Photobiol., 37, 587 (1983)

Wednesday

**Effects of Solvent on the Rate of Bacteriochlorophyll a Photooxidation**

Kenneth L. Marsh and John S. Connolly  
Photoconversion Research Branch  
Solar Energy Research Institute  
1617 Cole Boulevard  
Golden, CO 80401

Early investigations of the photostability of isolated bacteriochlorophyll a (BChl a) in organic media revealed significant solvent effects on the rate of pigment photooxidation. Thus, Smith and Calvin<sup>1</sup> reported a ~40-fold increase in the photodegradation rate of BChl a in acetone compared to the rate in diethyl ether. Recently, accurate values for the lifetimes ( $\tau_{\Delta}$ ) of singlet oxygen ( $O_2\ ^1\Delta_g$ ) have become available<sup>2</sup>, but these fail to account quantitatively for the observed solvent effect on photooxidation rate. Connolly et al.<sup>3</sup> estimated that the lowest triplet state of BChl a lies significantly lower than  $O_2\ ^1\Delta_g$  (7882  $cm^{-1}$ ) in polar or moderately polar solvents, and suggested that  $O_2$  may quench  $^3BChl\ a$  by an electron-transfer process which presumably would lead to degradation of the pigment. That  $O_2\ ^1\Delta_g$  is involved in photooxidation of BChl a appears certain, however, based on the cumulative evidence concerning photooxidation of other synthetic and natural magnesium porphyrins.<sup>4-7</sup> Moreover, Krasnovsky reported that BChl a is both a moderately effective sensitizer and a potent quencher of  $O_2\ ^1\Delta_g$  luminescence.<sup>8</sup>

The mechanistic aspects of BChl a photooxidation appear complicated based on the number (>5) of isolable photoproducts.<sup>9</sup> However, the spectrophotometric progress of the reaction in most solvents appears simple, exhibiting clear and well-defined isosbestic points that are preserved essentially to reaction completion, which indicates that the stoichiometry, however complex, remains constant.

We have carried out comparative studies of direct and rubrene-sensitized photooxidation of BChl a in nine solvents spanning a moderately wide range of dielectric constant (~2 to 37),  $O_2$ -solubility (~1-4 mM),  $\tau_{\Delta}$  (~16-100  $\mu s$ ) and solvent nucleophilicity; the last property affects the coordination state and the central Mg-ion of BChl a<sup>10</sup> and the fluorescence properties of the pigment.<sup>11</sup> The relative rates of pigment photodegradation show parallel trends in both the direct and sensitized experiments. However, there is no consistent correlation of initial photooxidation rates (in the direct irradiations) or relative degradation rates (compared to rubrene disappearance in the sensitized experiments) with either solvent polarity or  $\tau_{\Delta}$ . The only clear trend appears to be with the coordination number of BChl a magnesium. Thus, the rates are significantly slower when the central Mg is hexacoordinated (pyridine, p-dioxane, tetrahydrofuran, 2-methyltetrahydrofuran) compared to pentacoordinated Mg-species (acetonitrile, dichloromethane, benzene and acetone). It may be significant in this respect that the calculated fluorescence quantum yields<sup>11</sup> are higher when the Mg-coordination number is six, presumably reflecting lower intersystem crossing efficiencies. The sole exception to this trend is diethyl ether (pentacoordinated Mg) in which BChl a is remarkably stable.

A possible explanation of our results is that the triplet energy ( $E_T$ ) of BChl a lies close to the  $^1\Delta_g$  state of  $O_2$ <sup>12</sup> but varies with solvent.<sup>13</sup> Thus, in hexacoordinating solvents and in ether,  $E_T$  may be less than  $7882\text{ cm}^{-1}$ , but in pentacoordinating solvents (except ether)  $E_T > 7882\text{ cm}^{-1}$ . Experiments are currently in progress to verify this hypothesis; these include inhibition of direct photooxidation of BChl a by DABCO and  $\beta$ -carotene and laser flash photolysis experiments to determine the rates of quenching of  $^3\text{BChl a}$  by oxygen,  $\beta$ -carotene,<sup>3</sup> tetracene and pentacene.

#### References

1. J. R. Lindsay Smith and M. Calvin, J. Am. Chem. Soc. 88, 4500 (1966).
2. M. A. J. Rodgers, J. Am. Chem. Soc. 105, 6201 (1983).
3. J. S. Connolly, D. S. Gorman and G. R. Seely, Ann. N.Y. Acad. Sci. 206, 649 (1973).
4. J. Barrett, Nature, 215, 733 (1967).
5. J.-H. Fuhrhop, P. K. W. Wasser, J. Subramanian, and V. Schrader, J. Liebigs Ann. Chem. 1974, 1450-1466.
6. S. B. Brown, K. M. Smith, G. M. F. Bisset and R. F. Troxler, J. Biol. Chem. 255, 8063 (1980).
7. M. Krieg, G. S. Cox, and D. G. Whitten, XI International Conference on Photochemistry, College Park, Maryland, August 1983.
8. A. A. Krasnovsky, Jr., Photochem. Photobiol. 29, 29 (1979).
9. W. S. Kim, Biochem. Biophys. Acta, 112, 392 (1966).
10. T. A. Evans and J. J. Katz, Biochim. Biophys. Acta 396, 414 (1975).
11. J. S. Connolly, E. B. Samuel and A. F. Janzen, Photochem. Photobiol. 36, 565 (1983).
12. V. A. Shuvalov and W. W. Parson, Proc. Nat. Acad. Sci. USA 78, 957 (1981).
13. C. D. Tait and D. Holten, Photobiochem. Photobiophys. (in press, 1983).



Thursday

OXIDATION PHOTSENSITISED BY 2-CHLOROTHIOXANTHONE; THE ROLE  
OF SINGLET OXYGEN

R. B. CUNDALL, R.K. CHOWDHARY, A.K. DAVIES AND V.P. RUSSELL

Department of Biochemistry,

University of Salford,

Salford, M5 4WT. U.K.

2-Chlorothioxanthone (CTX) is a well known photoinitiator for the UV curing of polymer coatings. Its photoreactions are strongly inhibited by molecular oxygen which suggests that it might be an efficient photosensitiser by a singlet oxygen mechanism. CTX is practically insoluble in water but it can be incorporated into lipid vesicles and detergent micelles. Another interesting feature of this sensitiser is that its fluorescence is sensitive to the nature of the solvent. Thus, the fluorescence yield in non polar solvents is very low ( $\phi_f < 0.02$ ) whereas, in alcohols, an intense blue fluorescence is observed. This fluorescence is further enhanced by addition of water. It was hoped that CTX would perform the dual role of photosensitiser and fluorescence probe when incorporated into lipid vesicles and detergent micelles.

The photosensitised-oxidation experiments were carried out using a Rank Brothers oxygen electrode. The CTX was excited by light in the range  $\sim 380$  to  $\sim 430$  nm.

When CTX was solubilised in SDS micelles the solution exhibited an intense blue fluorescence. This indicates that the photosensitiser is not buried in the hydrophobic core of the micelles, but protrudes into the aqueous environment where hydrogen bonding to water molecules leads to fluorescence enhancement.

On addition of methionine to the CTX/SDS solution photosensitised absorption of oxygen occurred which, in the early stages, was linear. Addition of 1,4-diazabicyclo [2.2.2] octane (DABCO) quenched the oxygen uptake as did azide ion, consistent with a singlet oxygen-mediated oxidation of the methionine.

CTX was incorporated into the membranes of dipalmitoyl phosphatidylcholine (DPPC) vesicles. When an aqueous suspension of the vesicles was irradiated no oxygen absorption occurred. When vesicles were prepared from egg-yolk phosphatidyl choline rapid oxygen uptake occurred. In mixed vesicles, composed of DPPC and egg-yolk PC, the rate of photosensitised oxygen absorption increased with increasing concentration of egg-yolk PC.

The oxygen-uptake versus time traces for the CTX-photosensitised oxidation of egg PC differed significantly from those obtained for the CTX-photosensitised oxidation of methionine. After a period of 3 - 4 minutes, over which time the rate gradually increased, a high rate of oxygen uptake was observed which was almost linear - indicating the attainment of a steady state. When the light source was cut off, oxygen uptake continued in the dark, although at a much slower rate. The CTX-Photosensitised oxidation of egg yolk PC was inhibited by azide ion indicating the involvement of singlet oxygen. The rate of oxygen absorption was also reduced by the addition of p-methoxyphenol. This indicates that free radicals, as well as singlet oxygen, are involved in the CTX-photosensitised oxidation of egg-yolk PC vesicles.

SINGLET OXYGEN GENERATION FROM THE PEROXIDASE-CATALYZED AEROBIC OXIDATION OF  
AN ACTIVATED - CH<sub>2</sub>-SUBSTRATE

Nelson Durán<sup>S</sup>, Sandra T. Farias-Furtado<sup>S</sup>, Ana Campa<sup>E</sup>, Adelaide Faljoni-Alario<sup>E</sup>  
and Juan E. Brunet<sup>&</sup>

Instituto de Química, Biological Chemistry Laboratory, Universidade Estadual  
de Campinas, Cx.P. 6154, Campinas, SP, Brasil<sup>S</sup>, Universidade de São Paulo,  
Brazil<sup>E</sup> and Universidad Católica de Valparaíso, Chile<sup>&</sup>.

There is strong evidence that electronically excited products as well as those that give bioluminescence are present in biochemical systems<sup>(1-3)</sup>. Indirect evidence for this are: energy transfer processes, quenching, kinetics and comparison with photochemical processes<sup>(4,5)</sup>, and direct one is emission spectra<sup>(6,7)</sup>. Meanwhile peroxidase catalyzed aerobic oxidation of substrates such as indole-3-acetic acid<sup>(8)</sup>, 2-methylpropanal<sup>(5)</sup>, propanal<sup>(6)</sup>, acetoacetate<sup>(9)</sup> and many others generates triplet ketone in its excited states<sup>(1,2)</sup>, but there are few cases where the important emitting species seems to be singlet oxygen<sup>(7,10,11)</sup>.

Although we are aware of the limitation of biological systems in applying the techniques which have been developed using pure organic solvents<sup>(12)</sup>, methods which are nicely discussed by Foote<sup>(13)</sup>, we want to communicate a systematic study on the detection of singlet oxygen generation in peroxidase catalyzed oxidation of malonaldehyde (MA).

Correlation in O<sub>2</sub> consumption, product formation and Photon emission:

In the MA/HRP/O<sub>2</sub> system the oxygen consumption was much faster than light emission and product formation; this indicates accumulation of an intermediate, a fact which in turn suggests that the excited species may be formed outside the enzyme.

Conformational effect on the MA/HRP/O<sub>2</sub> Systems:

The variation of the spectra and its reactivity towards MA of solutions of HRP in dimethylsulfoxide has been studied. A broad pattern of changes was observed in the CD spectra of peroxidase, specially in the 400 nm region. These variations were correlated with conformational changes leading to a displacement of aromatic side chains in the heme environment<sup>(14)</sup>. Data has shown that the MA/HRP/O<sub>2</sub> system generates singlet oxygen probably outside the enzyme, i.e.,

in the bulk solutions. The actual data indicates that singlet oxygen production and its emission is completely insensitive to peroxidase conformational modifications. This is in agreement with the observation that the excited species in this system are formed outside the enzyme.

#### Photon emission: quenchers and enhancers:

The emission measured originated from dimol singlet oxygen was not affected by SOD, catalase, benzoate, bicarbonate, DB AS, but the emission was quenched by DOPA, histidine, methionine, tryptophan, guanosine and bilirubin, all excellent singlet oxygen quenchers. A small enhancement of the emission was observed in the presence of 10 mM DABCO. Also enhancement (4 fold) of the emission was observed in the presence of 10  $\mu$ M eosin and D<sub>2</sub>O (5 fold higher than in H<sub>2</sub>O). The effect of guanosine and histidine upon the emission of the MA/HRP/O<sub>2</sub> reaction was studied at concentration that did not influence the rate of oxygen consumption.

#### Products distribution:

Cadet and Teoule<sup>(17)</sup> reported the guanosine specifically react with singlet oxygen. In fact, using (5,8-<sup>14</sup>C) deoxyguanosine in the MA/HRP/O<sub>2</sub> system we detected N(2-deoxy-D-erythro-pentose) urea and N<sup>1</sup>(2-deoxy- $\beta$ -erythro-pentofuranosyl)N<sup>2</sup>-formylurea.

#### Emission spectrum:

In this system the intensity of the light was so intense, that it was possible to run the spectrum in a conventional fluorimeter in H<sub>2</sub>O or in D<sub>2</sub>O. The principal emission peak is just above 700 nm and occasionally a smaller peak was observed at 630 nm.

#### Kinetic studies:

Constancy of intercept in the plot of  $(I_0 - I_{A,Q})^{-1}$  versus  $(A)^{-1}$  at different (Q) is diagnostic for singlet oxygen quenching and implies that A and Q compete for a common intermediate. The quenching of histidine oxidation by guanosine in the MA/HRP/O<sub>2</sub> system gave constancy in the intercept. The ratio of reactivity of histidine to guanosine was ~5 in the photodynamic processes<sup>(16)</sup>; and in the ratio from Stern-Volmer plot of the enzymatic system gave a value of 4.3. A similar value was obtained from this steady state technique.

In conclusion, we suggest that the presence and role of singlet oxygen (<sup>1</sup> $\Delta$ g <sup>1</sup> $\Delta$ g) in the horseradish peroxidase-substrate system through photoenergized process can be ascertained in terms of combined data from photon emission, emission spectra, inhibition by quenchers, products analyses, D<sub>2</sub>O and DABCO effects, energy transfer to xanthenic dyes and kinetics analyses. This enzymatic system was proved to be very efficient for producing changes in ribosomes<sup>(11)</sup>, plasmid DNA<sup>(18)</sup> and in t-RNA<sup>(19)</sup>.

Support from FAPESP, CNPq, FINEP, UNESCO and OAS is ACKNOWLEDGED.

## REFERENCES

1. Cilento, G. (1980) Photochem. Photobiol. Rev. 5, 199-228.
2. Cilento, G. (1980) Acc. Chem. Res., 13, 225-230.
3. Durán, N., M. Haun and J.E. Brunet (1981) Rev. Latinoamer. Quim. 12, 1-11.
4. Durán, N. and G. Cilento (1980) Photochem. Photobiol., 32, 113-116.
5. Bechara, E.J.H., O.M.M.Faria-Oliveira, N. Durán, R.C. de Baptista and G. Cilento (1979) Photochem. Photobiol., 30, 101-110.
6. Haun, M., N. Durán, O. Augusto and G. Cilento (1980) Arch. Biochem. Biophys., 200, 245-252.
7. Vidigal-Martinelli, C., K. Zinner, B. Kachar, N. Durán and G. Cilento (1979) FEBS Lett., 108, 266-268.
8. De Mello, M.P., S.M. De Toledo, H. Aoyama, H.K. Sarkar, G. Cilento and N. Durán (1982) Photochem. Photobiol., 36, 21-24.
9. Takayama, K., M. Nakano, K. Zinner, C.C.C. Vidigal, N. Durán, Y. Shimizu and G. Cilento (1976) Arch. Biochem. Biophys., 176, 663-670.
10. De Mello, M.P., S.M. De Toledo, M. Haun, G. Cilento and N. Durán, (1980) Biochemistry, 19 5270-5275.
11. De Toledo, S.M., N. Durán and H. Singh (1983) Photobiol. Photobiophys 5, 237-243.
12. Durán, N. (1982) in Chemical and Biological Generation of Excited States (W. Adam and G. Cilento, Eds.) pp. 345-369, Academic Press Inc., New York.
13. Foote, C.S. (1979) in Biochemical and Clinical Aspects of Oxygen (W.S. Caughey, Ed.) pp 603-626, Academic Press Inc. New York.
14. Durán, N., J. Baeza, J. Freer, J.E. Brunet, G.A. Gonzalez, C.P. Sotomayor and A. Faljoni-Alário (1981) Biochem. Biophys. Res. Commun., 103, 131-138.
15. Foote, C.S. (1976) in Free Radicals in Biology (W.A. Pryor, Ed.) pp. 85-124, Academic Press Inc., New York.
16. Foote, C.S. (1978) in Singlet Oxygen, Reactions with Organic Compounds and Polymers (B. Branby and J.F. Rabek, Eds.) pp-135-146, John Wiley & Sons, New York.
17. Cadet, J. and R. Teoule (1978) Photochem. Photobiol., 28, 661-667.
18. De Toledo, S.M., A. Naha and N. Durán, (1982) Biochem. Biophys. Res. Commun., 104, 990-995.
19. Durán, N., M.C. Marcucci, M.P. De Mello and A. Faljoni-Alário (1983) Biochem. Biophys. Res. Commun., submitted.

## Singlet Oxygen Luminescence

B.M. Dzhagarov and K.I. Salokhiddinov

Institute of Physics, BSSR Academy of Sciences,  
Minsk 220602, USSR

A method is reported for the measurement of singlet molecular oxygen ( $^1\Delta_g$ ) lifetime in the solutions. The method is based on the pulsed laser excitation of the photosensitizing molecule and the  $^1\Delta_g \rightarrow ^3\Sigma_g^-$  luminescence registration with the help of the photodetector sensitive in the near IR spectral region. The developed method opens the way for studies of proper physico-chemical characteristics of singlet oxygen in different conditions, makes it possible to detect its presence and clarify its role in different photosensitization and oxidation processes.

Using this method or, as the need arose, also the method of laser nanosecond flash-photolysis, the following investigations have been made:

- (i) Lifetimes and relative luminescence yields of singlet oxygen in a variety of solvents have been measured.
- (ii) The mechanism of quenching the triplet states of compound organic molecules by molecular oxygen has been investigated, special emphasis being placed on tetrapyrrole molecules of different classes.
- (iii) A method has been proposed for measuring relative quantum yield of the intersystem crossing based on the comparison of singlet oxygen luminescence intensities, all other thing being equal. This method was used for the investigation of porphyrins and metalloporphyrins in order to study the intramolecular energetics parameters and select effective singlet oxygen sensitizers.

GAS PHASE GENERATION OF SINGLET OXYGEN  
AT ATMOSPHERIC PRESSURE BY DIRECT LASER EXCITATION

Walter C. Eisenberg, Alan Snelson, Ronald Butler and Kevin Taylor  
IIT Research Institute  
Chemistry and Chemical Engineering Research Division  
Chicago, Illinois

Robert W. Murray  
Department of Chemistry  
University of Missouri - St. Louis  
St. Louis, Missouri

The generation of singlet oxygen in the gas phase at atmospheric pressure was investigated using direct laser excitation. A Holobeam 255 Nd:YAG laser was used in the experiments. The output of the laser at 1065 nm lies within the  $O_2 \ ^3\Sigma_g (\nu=0) \rightarrow O_2 \ ^1\Delta_g (\nu=1)$  absorption band of the oxygen molecule at 1064 nm. Although the transition of ground state oxygen to the electronically excited singlet state is a "forbidden" process, we have previously shown that this transition does occur using simulated sunlight.<sup>1</sup>

Evidence for the formation of singlet oxygen by direct laser excitation was achieved in chemical trapping experiments with tetramethylethylene. These experiments were performed using a 12" by 1" OD pyrex tube having 1/8" quartz windows at each end. The tube containing the trapping agent and air or oxygen was placed in the beam of the laser during an experiment with the laser operating in a continuous mode. Control experiments involved no irradiation period. The production of detectable quantities of singlet oxygen was demonstrated in the conversion of tetramethylethylene to its known singlet oxygen product, 2,3-dimethyl-3-hydroperoxy-1-butene. No hydroperoxide was observed in the control experiments which were identical except there was no irradiation.

Further evidence for atmospheric gas phase generation of singlet delta oxygen by direct laser excitation was obtained by monitoring its characteristic 1.27  $\mu$ m emission using a germanium photodiode. In the second series of experiments, the laser was operated in a pulse mode. The laser beam was focused into an integrating sphere (excitation chamber) consisting of a 50 ml Pyrex

glass bulb with two 1/4" OD Kovar glass tubes attached at right angles. The surface of the integrating sphere was mirrored to permit many reflections of the laser beam. The optical detector was positioned at 90° with respect to the laser beam, and a light shutter was installed between the detector and the integrating sphere. The light shutter was open only when the laser was shut off. The laser was operated at full power for a period of 15 seconds, and was automatically shut off when the shutter was opened. The signal was led directly into an oscilloscope and data were collected only when oxygen was present in the integrating sphere. When nitrogen was used instead of oxygen or air, no 1.27  $\mu$ m emission was observed. The deactivation of singlet oxygen followed a first order decay with a mean value of  $k_T = 8.04 \text{ sec}^{-1}$  and a mean lifetime  $\tau$  of 0.124 seconds.

---

<sup>1</sup> Eisenberg, W. C., Taylor, K., Veltman, J., Murray, R. W., Generation of  $\text{O}_2 (^1\Delta_g)$  by Direct Absorption of Radiation by Ground State Oxygen, J. Am. Chem. Soc., 1982, 104, 1104-1105.



Wednesday

STUDIES ON SINGLET OXYGEN IN AQUEOUS SOLUTION

by D.F. Evans and M.W. Upton

Dept. of Chemistry, Imperial College, London SW7 2AY, England

The caesium and sodium salts of anthracene 9,10 bis ethane sulphonate (AES) are readily soluble in water over a wide range of pH, and react with  $^1\text{O}_2$  to give the endoperoxide as the sole detectable product. The reaction can be followed both from the U.V. absorption of AES and by HPLC of the endoperoxide. The trapping efficiency of AES for  $^1\text{O}_2$  has been measured as a function of concentration. AES does not react with  $\text{O}_2^-$  or  $\text{HO}_2$ , but acts as a trap for radicals such as  $\cdot\text{OH}$ . The products, however, can readily be distinguished from the endoperoxide by HPLC.

AES has been used to study quantitatively the production of singlet oxygen in a variety of reactions occurring in aqueous solution.

(1) Two electron oxidation of  $\text{H}_2\text{O}_2$ .

The reaction of  $\text{H}_2\text{O}_2$  with N chlorosuccinimide and Chloramine T produces  $^1\text{O}_2$  with  $\sim 100\%$  efficiency. With Chloramine T, the reactive species is dichloramine T produced by disproportionation.

Both iodoxybenzene and iodosobenzene sulphonic acid react with  $\text{H}_2\text{O}_2$  to give oxygen, of which  $\sim 45\%$  only is in the singlet state. This may be due to spin-orbit coupling involving the heavy iodine atom. With  $\text{IO}_4^-$  and  $\text{H}_2\text{O}_2$ , the yields of  $^1\text{O}_2$  are irreproducible, and slightly lower, indicating a concomitant free radical reaction.

(2) Decomposition of peroxyacids at their pK. When sufficient precautions to avoid catalysis by transition metal ions are taken, the decomposition of peroxymonophthalic, peroxyacetic and peroxymono sulphuric acids give  $^1\text{O}_2$  in essentially quantitative yields. The reported reaction of  $\text{H}_2\text{O}_2$  with peroxyacid anions does not produce  $^1\text{O}_2$ , and in the absence of metal catalysis is very slow indeed, if it occurs at all.

(3) Catalyzed decomposition of  $\text{H}_2\text{O}_2$ .

Friday

$O_2(a^1\Delta_g)$  SENSITIZED CHEMILUMINESCENCE OF  $a^1\Delta \rightarrow X^3\Sigma^-$  AND  
 $b^1\Sigma^+ \rightarrow X^3\Sigma^-$  TRANSITIONS OF GROUP VI-VI AND GROUP V-VII

DIATOMIC MOLECULES

R. Winter, H. Kruse, E. H. Fink, J. Wildt and F. Zabel  
Physikalische Chemie - FB 9, Bergische Universität - GH Wuppertal  
D-5600 Wuppertal 1, West Germany

Abstract

Systematic studies of the near infrared emissions from the  $b^1\Sigma^+$  and  $a^1\Delta$  states of group VI-VI and group V-VII diatomic molecules in chemiluminescence systems, following the excimer laser photolysis of suitable parent compounds, and by Raman-shifted dye laser fluorescence (LIF) are reported. The  $b \rightarrow X$  band systems have been observed for 25 molecules of the two groups mostly by studying the  $O_2(a^1\Delta_g)$  sensitized chemiluminescence in fast flow systems. For SO also the (0,0) band of the  $a \rightarrow X$  transition has been detected. Using a Fourier-Transform-IR-Spectrometer, rotationally resolved spectra of the  $b \rightarrow X$  bands of  $S_2$  and SeS were measured. Time resolved studies of the  $b \rightarrow X$  emissions by excimer laser photolysis and LIF yielded preliminary results for the radiative lifetimes and the quenching rate constants of the b states of SO, SeO, PCl, PBr, and SbCl. These studies as well as Stern-Volmer type measurements in flow systems strongly indicate that the quenching of the b states in many cases proceeds by near-resonant  $E \rightarrow V$  energy transfer. As will be shown, a wealth of other energy transfer and energy pooling processes is operative in the chemiluminescence and photolysis systems.

Thursday

## SINGLET OXYGEN PRODUCTION FROM PHOTODYNAMIC SENSITIZERS

C. S. Foote and D. C. Dobrowolski

Department of Chemistry and Biochemistry

University of California, Los Angeles

Los Angeles, California 90024

There is a wide range of naturally occurring photodynamically active compounds from plant sources. Examples include the fungal pigment cercosporin and a variety of plant polyacetylene derivatives. These compounds probably serve to protect the synthesizing organism from animal (mammal or insect) browsing, or, in the case of cercosporin, to promote the attack of the cercospora fungus on the target cell. Another interesting compound is 4-thiouridine, a nucleoside found in transfer RNA of certain wild-type e. coli strains. This nucleoside appears to be the sensitizer in an oxygen-sensitive near-UV photodynamic single strand breakage of DNA in these organisms.

Using the 1268 nm emission from singlet oxygen, it is relatively simple to quantitate the production of singlet oxygen from many sensitizers. The sensitizer is excited by a short pulse from a Nd:YAG laser, and the weak luminescence from singlet oxygen is detected in a time resolved system. The intensity of the luminescence is compared with that produced from a sensitizer with a known singlet oxygen quantum yield. These quantum yields are checked by measuring the quantum yield of photooxidation of typical singlet oxygen acceptors; the results of the

control experiments agree well with those of the luminescence experiments in all cases so far studied. Using this technique, cercosporin, terthienyl, and 4-thiouridine have all been found to produce singlet oxygen in solution in substantial yields.

The efficient production of singlet oxygen from these sensitizers is a necessary but not sufficient condition for the intermediacy of singlet oxygen in the photodynamic damage sensitized by these compounds. That is, singlet oxygen production could be only a side reaction accompanying the actual photodynamic processes. The strong protective effect of singlet oxygen quenchers against damage mediated by cercosporin and terthienyl makes it nearly certain that singlet oxygen is the active species with these pigments. However, further work will be required for the 4-thiouridine case, since the effects of inhibitors do not seem to be totally explicable on the basis of a pure singlet oxygen mechanism.

Wednesday

## ASPECTS OF PHOTSENSITIZED LIGNIN OXYGENATION

Ingegerd Forsskåhl

Department of Chemistry, University of Helsinki,  
Et. Hesperiank. 4, 00100 Helsinki 10, Finland

Substantial knowledge has been accumulated of the chromophoric structures in native lignin and in modified degraded lignin. These structures, acting as sensitizers, together with light and oxygen are directly or indirectly responsible for reactions leading ultimately to photo-degradation of the starting material. This is often displayed by the darkening or the yellowing of wood specimen and industrial articles made of wood, such as high yield masses and different paper products. It has been of interest for the economical and ecological development to reduce the use of bleaching chemicals and to prevent re-yellowing of the products.

Earlier studies have revealed the emphasis of oxygen in photochemical lignin degradation and that the process involves oxygenation pathways.

This work is a contribution to the understanding of the mechanism of the yellowing process and clearly shows the complexity of the reaction.

The results are derived from model experiments with tert-butylguaiacylcarbinol sensitized by aromatic  $\alpha$ -carbonyl triplets and by compounds containing ring-conjugated double bonds. The rates of disappearance of the reactants were determined by gas chromatography and the formation of peroxides were measured by iodometric titration. The kinetics of the reactions were also studied in the presence of "so called singlet oxygen quenchers and free radical scavengers.

The results indicate that the mechanism for photo-oxygenation is very complicated with many pathways operating simultaneously, one of which is attributed to a reaction

involving singlet oxygen in addition to the known hydrogen abstraction process. Radical reactions including thermo- or photochemically instable peroxides are also of significance, particularly during prolonged irradiations.

The relative importance of these pathways is discussed and the course of the reaction depends largely on the conditions, such as the concentrations, the properties of the reactants and the reaction time. This means that one mechanism can dominate in one system, whereas another is of major importance in another one.

THE EFFECT OF STRAIN ON THE REARRANGEMENTS  
OF PRIMARY SINGLET OXYGEN PRODUCTS

ARYEH A. FRIMER

Department of Chemistry, Bar-Ilan University, Ramat-Gan 52100, Israel

We have argued<sup>1</sup> previously that  $^1\text{O}_2$  is relatively insensitive to strain consideration either present in the starting material or developing in the product. More important factors seem to be the ground state geometry of the olefin, the  $\text{C}_\alpha - \text{H}_{\text{allylic}}$  interatomic distance and the ionization potential of the double bond. Strain does, however, play a crucial and fascinating role in the secondary rearrangements of primary  $^1\text{O}_2$  products. The plethora of products obtained can be easily rationalized in terms of intriguing variations on the "Hock-cleavage" theme.<sup>2</sup> These include ring rather than vinyl group migration and divinyl ether formation in competition with hemiacetal production yielding oxetanes, oxetenes and alkylidene oxetenes as intermediates.

<sup>1</sup> A.A. Frimer, Israel J. Chem., 21, 194 (1981).

<sup>2</sup> A.A. Frimer, Chem. Rev., 79, 359 (1979).

**DYE-SENSITIZED PHOTO-OXYGENATION OF ISONIAZID  
BY SINGLET MOLECULAR OXYGEN**

RAKESH D. DUBEY\* & P.B. GANDHI  
SCHOOL OF STUDIES IN CHEMISTRY,  
VIKRAM UNIVERSITY,  
UJJAIN - 454 010 (M.P.)  
INDIA.

**A B S T R A C T**

The present paper deals with the formation of a dicarbonyl cyclic compound from the isoniazid, by the dye-sensitized photo-oxygenation method. The formation of product has been explained by both Gollnick's "type I" and "type II" reaction mechanisms. The participation of singlet oxygen has been confirmed by the use of different dyes and quenchers.

The structure of the product has been proposed on the basis of electronic absorption, infrared, nmr and mass spectral data.

The yield of product in different conditions has been also determined.

---

\* Present Address  
ENVIRONMENTAL PLANNING &  
COORDINATION ORGANISATION  
E-5 SECTOR ARERA COLONY  
BHOPAL - 462016 (M.P.)  
INDIA



Wednesday

The Efficiency of Triplet Energy Transfer to  $O_2(^3\Sigma_g^-)$ :

Sensitizer and Medium Dependence

A. A. Gorman<sup>1</sup>, I. Hamblett<sup>1</sup> and M. A. J. Rodgers<sup>2</sup>




Chemistry Department, University of Manchester, Manchester M13 9PL<sup>1</sup>

and

The Center for Fast Kinetics Research, University of Texas at Austin,

Austin, Texas 78712<sup>2</sup>

The efficiency of formation of  $O_2(^1\Delta_g)$  in the process whereby electronically excited triplet states are quenched by  $O_2(^3\Sigma_g^-)$  has been a matter of some debate. The two spin-allowed processes are represented by channels 1 and 2 which proceed *via* the singlet and triplet manifolds



respectively of the encounter complex. Quenching *via* the quintet manifold is forbidden. Rate constants for oxygen quenching of triplet states do not *generally* exceed  $\sim 1/9$ th of the diffusion controlled value and it was naturally concluded that quenching is restricted to the singlet manifold and that each quenching act leads to a molecule of  $O_2(^1\Delta_g)$ .<sup>1</sup> The validity of this conclusion has been seriously questioned as a result of pulse radiolysis<sup>2</sup> and pulsed laser<sup>3</sup> studies which in turn have themselves been the subjects of criticism based on steady-state experiments.<sup>4</sup>

/over.....

- 2 -

In the work to be described a series of sensitizers in aerated media have been subjected to pulsed laser excitation at 355 nm. In each case the sensitizer triplet yield and the  $O_2(^1\Delta_g)$  emission intensity have been monitored by kinetic absorption and kinetic emission spectroscopy respectively. The significance of these data will be discussed.

#### References


1. O. L. J. Gijzeman, F. Kaufman and G. Porter, J. Chem. Soc., Faraday Trans.2, 69, 708 (1973).
2. A. A. Gorman, G. Lovering and M. A. J. Rodgers, J. Amer. Chem.Soc., 100, 4527 (1978).
3. A. Garner and F. Wilkinson, "Singlet Oxygen - Reactions with Organic Compounds and Polymers", B. Ranby and J. F. Rabek, eds., Wiley, New York, 1978, p.48.
4. B. Stevens, K. L. Marsh and J. A. Barltrop, J.Phys. Chem., 85, 3079 (1981).

Wednesday

## SENSITIZED PHOTOOXIDATION OF METHYL SUBSTITUTED CYCLOBUTENES

Gary W. Griffin, Gary P. Kirschenheuter and Pesi P. Umrigar  
University of New Orleans


The series of methyl-substituted 1,2-diphenylcyclobutenes 1a-d were subjected to photooxidation in the presence of sensitizers including methylene blue (MB) and rose bengal (RB).



- 1a  $R^1=R^2=R^3=R^4=Me$   
1b  $R^1=R^2=R^3=Me, R^4=H$   
1c  $R^1=R^3=Me, R^2=R^4=H$  (trans)  
1d  $R^1=Me, R^2=R^3=R^4=H$

Sensitized photooxidations of 1a with MB and RB lead to formation of the ring-contracted product 6 presumably by initial "4+2" addition of singlet oxygen to the styryl diene moiety of the substrate.

When the electron-transfer sensitizer 9,10-dicyanoanthracene is used, the reactions of 1 take a markedly different course. For example, 1a is rapidly photooxidized in  $CH_3CN$  to give the spectrum of products 2-5 through a mechanism in which superoxide radical anion is implicated. Thus tetramethylcyclobutene 1a may be employed to differentiate between photo-oxygenations involving singlet oxygen and superoxide ion. The selectivity exhibited by this substrate has been used to probe the identity of the cytotoxic agent responsible for the dramatic cancer cures observed with hematoporphyrin derivative (HPD) as a phototherapeutic agent. Our preliminary data using HPD as a sensitizer *in vitro* supports Dougherty's contention that singlet oxygen is involved although his conclusion was based on trapping experiments with 1,3-diphenylisobenzofuran, an assay which must now be considered questionable.



### Reactions of Singlet Oxygen with Liposomes

Leonard I. Grossweiner

Biophysics Laboratory, Physics Department, Illinois Institute of Technology, Chicago, IL 60616, U.S.A.

Liposomes have been used as model membranes for the investigation of chemical and biological processes photosensitized by natural and synthetic pigments. Work in this laboratory has been done with egg phosphatidylcholine (EPC) liposomes photosensitized by methylene blue (MB) <sup>1,2</sup>, riboflavin (RF)/EDTA <sup>1</sup>, hematoporphyrin (HP) <sup>3</sup>, the constituents of hematoporphyrin derivative (HPD) <sup>4-6</sup>, and 8-methoxypsoralen (8-MOP) <sup>7</sup>. Damage sensitized by these agents, incorporated in the liposomes and added to the external aqueous phase, has been assayed by membrane lysis, marker release, and lipid peroxidation. The involvement of singlet oxygen (<sup>1</sup>O<sub>2</sub>) was demonstrated for MB, HPD, dilute HP and 8-MOP; aggregated HP photosensitized via an anaerobic, Type I mechanism; superoxide (O<sub>2</sub><sup>-</sup>) was implicated for RF/EDTA. Dipalmitoylphosphatidylcholine liposomes were not sensitized by these agents, implicating the unsaturated fatty acids as the site of <sup>1</sup>O<sub>2</sub> and O<sub>2</sub><sup>-</sup> attack. Liposome damage was inhibited by water soluble acceptors, azide ion, DABCO, and human serum albumin, indicating that <sup>1</sup>O<sub>2</sub> generated in the liposomes diffused into the external medium prior to reacting with the membrane. The efficiency was estimated by generating <sup>1</sup>O<sub>2</sub> in the gas phase and bubbling into the liposome suspension. The analysis based on large target diffusion kinetics indicates that 3000 <sup>1</sup>O<sub>2</sub> interactions at the liposome surface were required for lysis of 1.5 μm EPC liposomes <sup>8</sup>. The diffusion of HP and HPD into sonicated EPC liposomes leads to a large red shift of the Soret bands and enhanced fluorescence efficiency, indicative of monomerization. Visible light irradiation of the dyed liposomes induced lipid peroxidation, with positive mechanistic tests for <sup>1</sup>O<sub>2</sub>. The occurrence of these effects with the tumor localizing and photosensitizing constituent of HPD strongly suggests that the in situ tumor sensitizer in photoradiation therapy is a monomerized, fluorescent constituent of HPD localized in a lipophilic environment that generates <sup>1</sup>O<sub>2</sub> when excited by visible radiation.

L. I. Grossweiner

This work was supported by NIH Grant GM 20117 and Department of Energy Contract No. AC02-76EV02217.

References

1. R. Muller-Runkel, J. Blais and L. I. Grossweiner, Photochem. Photobiol. 33, 683 - 687 (1981).
2. L. I. Grossweiner and J. B. Grossweiner, Photochem. Photobiol. 35, 583 - 586 (1982).
3. L. I. Grossweiner, A. S. Patel and J. B. Grossweiner, Photochem. Photobiol. 36, 159 - 167 (1982).
4. L. I. Grossweiner and G. C. Goyal, Photochem. Photobiol., 37, 529 - 532 (1983).
5. L. I. Grossweiner, The Clayton Foundation Symposium on Porphyrin Localization and Treatment of Tumors, Santa Barbara, CA, April 24 - 28, 1983.
6. G. C. Goyal, A. Blum and L. I. Grossweiner, Cancer Res. xx, xxx - xxx (1983).
7. R. Muller-Runkel, Ph.D. Thesis, Illinois Institute of Technology, May, 1981.
8. W. C. Eisenberg, K. Taylor and L. I. Grossweiner, Photochem. Photobiol. (submitted).

PROSPECTS FOR SOLAR SYNTHESIS. I. A NEW METHOD FOR SINGLET OXYGEN REACTIONS USING NATURAL SUNLIGHT. J. E. Guillet, J. Sherren, H. M. Gharapetian and K. MacInnis. Department of Chemistry, University of Toronto, Toronto, Canada M5S 1A1.

The study of organic photochemistry has provided a variety of interesting synthetic routes to novel organic compounds and intermediates. Unfortunately, the high cost of using electrically-driven lamps to provide the necessary photon energy has made photochemical steps impractical in most commercial applications. We wish to report a new method of carrying out organic photochemical reactions using sunlight as the photon source, which is both inexpensive and easily scaled up for commercial synthesis.

In this procedure, the organic reagent is absorbed in transparent beads of crosslinked ethylene-vinyl acetate copolymer. These beads float readily on water, and can be exposed to natural sunlight on the surface of a small pond or pool. After exposure, the beads are collected, washed with water, and dried in air. The products and unreacted starting material are removed by selective extraction with suitable solvent mixtures, after which the beads can be recycled through the process. The photoreaction can be carried out either batchwise or in a continuous process. In the dry form the beads can be moved by airveyor, while during exposure they are most conveniently moved by adjusting the flow of water through the pond-reactor.

A number of classical photochemical reactions have been studied in these systems, including photoisomerizations, the Norrish type II reaction in various ketones, and the addition of singlet oxygen to olefinic double bonds. The latter process is of particular interest, because it is sensitized by compounds absorbing visible light near the maximum intensity of the solar spectrum. Furthermore, the beads have a relatively high permeability to oxygen from the air.

In the present work the rates of addition of singlet oxygen to squalene were measured. Rose Bengal was used as the sensitizer. The resulting allylic hydroperoxide was estimated by the triphenyl phosphine method. Conversions of up to 10% were obtained after four days exposure to sunlight. The results of singlet oxygen additions were compared with those for the type II reaction of aromatic ketones.

Mechanism of Singlet Oxygen Luminescence  
Sensitization by Polyatomic Molecules

G.P. Gurinovich

Institute of Physics of the BSSR Academy of Sciences, Minsk  
220602, USSR

Of the various methods of singlet oxygen ( $^1O_2$ ) generation, the formation of  $^1O_2$  by dye- and pigment sensitization is of particular interest both for studying different quenching mechanisms of organic molecule excited (singlet and triplet) states and for understanding chemical, biological and medical light-initiated processes involving oxygen.

This paper presents a thorough consideration of mechanisms of quenching the excited states of the sensitizer's molecules by oxygen resulting in the formation of  $^1O_2(^1\Delta_g)$ , primary attention being focused on utilizing the  $^1O_2(^1\Delta_g)$  own luminescence for dealing with the following problems:

- oxygen quenching of sensitizer fluorescence with different  $E(S_1-T_1)$  energy intervals;
- quenching of the sensitizer molecules' triplet states;
- intermolecular energy transfer involving oxygen;
- $^1O_2(^1\Delta_g)$  luminescence resulting from interaction of oxygen molecules with excited molecules of the sensitizer;
- photosensitized  $^1O_2(^1\Delta_g)$  reactions in solutions.

Friday

BEHAVIOR OF SINGLET OXYGEN IN THE  
OXYGEN-IODINE TRANSFER LASER


R. F. Heidner III

Aerophysics Laboratory  
The Aerospace Corporation  
Los Angeles, CA 90009

ABSTRACT

$O_2(^1\Delta)$  supplies all the energy required to produce the laser medium in the chemical oxygen-iodine (COIL) laser. Since  $O_2(^1\Delta)$  can be made by the chemical reaction between  $Cl_2$  and basic hydrogen peroxide, this near-infrared laser ( $\lambda = 1.315 \mu m$ ) is one of the very few all-chemical laser devices.

It is necessary to produce high densities of  $O_2(^1\Delta)$ , however, it is equally important that one produce high ratios of  $O_2(^1\Delta)/O_2(^3\Sigma)$ . The electronic state equilibrium



$$K_{eq} = 2.9 \quad (T = 295K)$$

will support gain on the  $I(^2P_{1/2} \rightarrow ^2P_{3/2})$  transition ( $I^*/I > 0.5$ ) only if  $O_2(^1\Delta)/O_2(^3\Sigma) > 0.17$ . Thus, the loss processes for  $O_2(^1\Delta)$  are critical for the energetic efficiency of this laser. This talk will review the loss processes associated with the presence of atomic and molecular iodine. These may be summarized as follows:



- (1)  $I_2$  dissociation by  $O_2(^1\Delta)$
- (2)  $I^*$  pumping by  $O_2(^1\Delta)$
- (3) Quenching of  $O_2(^1\Delta)$  by I-atom related processes

These processes will be considered in the context of the conventional cw COIL laser system. In addition, recent experimental work on the kinetics of a premixed, pulsed version of this promising laser system will be presented.

Friday

Laser Kinetic Studies of  $I(^2P_{1/2})$  and  $O_2(^1\Delta_g)$ 

G. E. Hall, W. Marinelli, Sivaram Arepalli,

A. T. Young, P. L. Houston, and J. R. Wiesenfeld

Department of Chemistry  
 Cornell University  
 Ithaca, New York 14853

Multiphoton ionization has been used to detect sensitively  $O_2(^1\Delta)$ , both in a microwave discharge and from the 266-nm photolysis of  $O_3$ . The spectra showed several vibrational levels for each of two Rydberg transitions in the 260-245 nm region as well as a valence band near 340 nm. None of the  $O_2(^1\Delta)$  states has previously been reported. Their assignment is currently in progress.

The equilibrium between excited iodine atoms and oxygen has been examined by monitoring the time-dependent  $I^*[\equiv I(^2P_{1/2})]$  concentration following creation of this species by pulsed laser photolysis of iodine-containing precursors. A double exponential decay is observed which reflects a fast approach to equilibrium followed by a slower relaxation of the equilibrated mixture. Data obtained with the  $I^*$  precursors HI and  $CH_3I$  show that the rate constant for the reaction  $I^* + O_2 \rightarrow I + O_2$  is insignificant compared to that for the reaction  $I^* + O_2 \rightarrow I + O_2(^1\Delta)$ . Data obtained from the precursors  $i-C_3F_7I$ ,  $n-C_3F_7I$ ,  $C_2H_5I$ , and  $CF_3I$  suggest that  $O_2(^1\Delta)$  is rapidly relaxed by the precursor itself, by the precursor radical, or by some product formed in a reaction between the precursor radical and oxygen. The rate constant for the process  $I^* + O_2 \rightarrow I + O_2(^1\Delta)$  has been found to be  $(8.8 \pm 0.9) \times 10^5 \text{ sec}^{-1} \text{ torr}^{-1}$ , while that for the process  $I^* + O_2 \rightarrow I + O_2$  is  $(0.3 \pm 1.3) \times 10^5 \text{ sec}^{-1} \text{ torr}^{-1}$ .

Electronic-to-vibrational energy transfer from  $I^*(5^2P_{1/2})$  to  $I_2(25 < v < 43)$  has been observed.  $I^*$  was created by pulsed laser photolysis of either  $I_2/Ar$  mixtures at 475 nm or  $CF_3I/I_2/Ar$  mixtures at 266 nm, while the resulting vibrational distribution of  $I_2$  was monitored by laser induced fluorescence on the  $I_2(B+X)$  transition. The experimental results are consistent with a nascent  $I_2$  product distribution which is inverted, with a substantial fraction of the  $I_2$  molecules formed in  $v > 30$ . Roughly 2% of the  $I^*$  deactivations result in  $I_2(v=40)$ . The rate constants for vibrational relaxation of  $I_2(v=40)$  by argon, helium, and  $I_2$  at room temperature are  $(7.3 \pm 0.3) \times 10^5 \text{ sec}^{-1} \text{ Torr}^{-1}$ ,  $(1.0 \pm 0.2) \times 10^6 \text{ sec}^{-1} \text{ Torr}^{-1}$ , and  $(1.8 \pm 0.4) \times 10^6 \text{ sec}^{-1} \text{ Torr}^{-1}$ , respectively. These results have important implications for the mechanism of  $I_2$  dissociation in the chemical oxygen/iodine laser. A chain branching mechanism consisting of the steps  $I^* + I_2 \rightarrow I + I_2(20 < v < 40)$ ,  $I_2(20 < v < 40) + O_2(^1\Delta) \rightarrow 2I + O_2$ , and  $O_2(^1\Delta) + I \rightarrow O_2 + I^*$  may be responsible for the dissociation.

Saturday

Construction of 1,2,4-Trioxane rings by Photo-oxygenation of Enol Ethers

Charles W. Jefford, John Boukouvalas and Shigeo Kohmoto

Department of Organic Chemistry, University of Geneva

1211 Geneva 4, Switzerland

Enol ethers which lack the option of forming hydroperoxides react readily with singlet oxygen to give 1,2-dioxetanes by a mechanism which could involve charge-transfer complexes, radical ion pairs, biradical or zwitterionic species. We report that dioxetane ring formation can be intercepted when aldehydes are used as solvents. They are efficiently trapped undergoing conversion to 1,2,4-trioxanes, thereby providing proof for an intermediate zwitterion.

In other cases, 1,2-dioxetanes and 1,4-endoperoxides, products typically obtained by photo-oxygenation, can be opened by acid catalysis to give the corresponding stable  $\beta$ -hydroperoxy cations which are either sufficiently long-lived or reactive enough towards carbonyl compounds to give trioxanes. The mechanistic and synthetic aspects of these new reactions will be discussed.

In view of the paucity of structural information on 1,2,4-trioxanes, illustrations of conformational features will be given in the light of recently obtained X-ray analytical and NMR data.


SINGLET OXYGEN PRODUCTION BY LACTOPEROXIDASE:  
HALIDE DEPENDENCE AND QUANTITATION OF YIELD

Jeffrey R. Kanofsky

Edward Hines, Jr., VA Hospital, Hines, IL

Preliminary studies in this laboratory have demonstrated singlet oxygen (SO) production in the lactoperoxidase (LPO)-H<sub>2</sub>O<sub>2</sub>-Br<sup>-</sup> system using 1268 nm chemiluminescence (CL) (JBC 258: 5991, 1983). Further details about the technique and the LPO-H<sub>2</sub>O<sub>2</sub>-halide systems are now reported. For the study of biochemical systems, 1268 nm CL is both more sensitive and more selective than dimole CL. The current CL spectrometer can measure SO production rates of 30 nM/min (flow system study using the H<sub>2</sub>O<sub>2</sub> + NaOCl reaction as the SO source). In systems where D<sub>2</sub>O solvent can be used, this decreases to 1 nM/min. In contrast, the sensitivity of dimole CL (cooled RCA 8852 photomultiplier using single photon counting) is 1000 nM/min. Of more significance is the empirical observation that CL at 1268 nm from sources other than SO is rare, while biologic oxidations often produce CL at visible and near infrared wavelengths including the dimole bands. For example, HOBr, an intermediate in the LPO-H<sub>2</sub>O<sub>2</sub>-Br<sup>-</sup> system, reacts with albumin to produce CL with a broad spectral distribution overlying the dimole bands, but no detectable CL at 1268 nm. However, there is no reason to think that 1268 nm CL is totally specific for SO in complex systems, so additional supporting evidence to confirm SO production should be obtained. This includes (1) spectral analysis, (2) a large isotope effect in D<sub>2</sub>O of about 30 and (3) inhibition of CL with appropriate SO quenchers and traps (Quenching is not highly specific for SO, but the failure to quench is strong evidence against SO.).

Under optimal conditions (0.1 mg/ml LPO, 3 mM NaBr, pH 4.5 and 1 mM H<sub>2</sub>O<sub>2</sub>) the LPO-H<sub>2</sub>O<sub>2</sub>-Br system stoichiometrically produces SO by the mechanism



The bromine containing species HOBr, Br<sub>2</sub> and Br<sub>3</sub><sup>-</sup> are in rapid equilibrium, so it is not possible to determine which of these is the initial product of the LPO catalyzed oxidation of Br<sup>-</sup>. While oxidized bromine species are important reaction intermediates, they are quantitatively minor final system products.

## Kanofsky-2

LPO does not produce SO with either  $\text{Cl}^-$  or  $\text{I}^-$  as a substrate.  $\text{Cl}^-$  is not oxidized. Past studies have shown that  $\text{I}^-$  is quantitatively converted to  $\text{I}_2$  (Anal Biochem 42: 195, 1971).

Thursday

ENZYME GENERATION OF SINGLET ( $^1\Delta_g$ ) MOLECULAR OXYGEN:  
NEAR IR SOLUTION EMISSION SPECTROSCOPY

Ahsan U. Khan  
Institute of Molecular Biophysics and Department of Chemistry  
Florida State University  
Tallahassee, Florida 32306

## ABSTRACT

Enzyme production of singlet ( $^1\Delta_g$ ) molecular oxygen in a dark reaction is revealed via near IR emission spectroscopy in solution. Singlet molecular oxygen emission [(0,0)  $^1\Delta_g \rightarrow ^3\Sigma_g^-$  at 1.28 micron] is observed in the enzyme systems (1,2): chloroperoxidase/H<sub>2</sub>O<sub>2</sub>/Cl<sup>-</sup> and lactoperoxidase/H<sub>2</sub>O<sub>2</sub>/Br<sup>-</sup>. In the catalase/H<sub>2</sub>O<sub>2</sub> system a strong emission at 1.64 micron with bandwidth of 753 cm<sup>-1</sup> is observed, tentatively interpreted as arising from perturbed  $^1\Delta_g$  state of molecular oxygen.

The development of near IR solution emission spectroscopy has enabled several other seminal observations: (i) the generation of O<sub>2</sub> ( $^1\Delta_g$ ) in the disproportionation of the superoxide anion, O<sub>2</sub><sup>-</sup> (3); (ii) quenching of O<sub>2</sub> ( $^1\Delta_g$ ) by vitamin C in aqueous solution (4); and (iii) solvent perturbed O<sub>2</sub> ( $^1\Delta_g$ ) emission (5).

1. A. U. Khan, P. Gebauer and L. P. Hager, Proc. Natl. Acad. Sci. USA, 80, 5195 (1983).
2. A. U. Khan, J. Am. Chem. Soc., to appear Nov. 30, 1983
3. A. U. Khan, J. Am. Chem. Soc. 103, 6516 (1981).
4. P. Chou and A. U. Khan, Biochem. Biophys. Res. Commun., 115, 932 (1983).
5. P. Chou and A. U. Khan, accepted in Chem. Phys. Letters, 1983.

---

Work supported by the National Foundation for Cancer Research, Bethesda, MD.

# PHOTOSENSITIZED LUMINESCENCE OF SINGLET OXYGEN IN SOLUTIONS: MECHANISMS OF QUENCHING BY SOLVENTS AND BIOLOGICALLY IMPORTANT COMPOUNDS.

A.A. Krasnovsky Jr.

Department of Biology, Moscow State University, Moscow, USSR.

The photosensitized luminescence of singlet oxygen in pigment solutions discovered in our experiments several years ago / I / is now intensively used as a tool for investigation of  $^1\Delta_g$  state of molecular oxygen. In many cases this tool yields more reliable data than methods based on the use of  $^1\Delta_g$  acceptors. The present paper summarizes the results of our measurements of the rate constants of  $^1\Delta_g$  quenching ( $K_q$ ) made with the aid of the oxygen luminescence / I, 2 and refs. therein /.

It has been found that  $K_q$  values of n-alkanes (pentane-dodecane) increase with the increase of the number of hydrogen atoms ( $n_H$ ) in alkane molecules. The  $K_q/n_H$  ratio is constant ( $\sim 450 \text{ M}^{-1}\text{s}^{-1}$ ). It can be regarded as an activity of a single CH group. For n-alcohols (methanol-decanol) and saturated organic acids (acetic-behenic) this ratio declines with the increase of  $n_H$ . Analysis leads to conclusion that the activities of CH and OH groups are different, OH groups being more active by an order of magnitude. Using the experimental values of  $K_q$  it is shown that  $^1\Delta_g$  lifetimes ( $\tau_\Delta$ ) and probabilities of light emission by the  $^1\Delta_g$  state ( $p_r$ ) in all the solvents studied are well approximated by the following equations:

$$\tau_\Delta = 1/(K_q[S]); \quad p_r = 1/(K_q\tau_r[S])$$

[S] is the molar concentration of solvents;  $\tau_r$  is the radiative lifetime of  $^1\Delta_g$  equal to 4 s.

The energy transfer from  $^1\Delta_g$  to vibrational sublevels of organic molecules is the most probable mechanism of  $^1\Delta_g$  quenching by the mentioned compounds. The same mechanism determines the activities of saturated fatty acids and lipids, while its contribution to the overall quenching effect of unsaturated fatty acids varies from 40 to 45 depending on the number of double bonds in fatty acid molecules. Chemical oxygenation of these molecules appears to play a more important role in their interaction with  $^1\Delta_g$ .

Chlorophylls and porphyrins were found to be efficient physical quenchers of  $^1\Delta_g$ . Their activities depend on the macrocycle structure as well as on the presence and nature of the central metal atom. Two mechanisms appear to determine the quenching effect of these compounds - energy transfer from  $^1\Delta_g$  to low lying electronic levels of quenchers or formation of exciplexes with charge transfer between  $^1\Delta_g$  and porphyrins. The first mechanism is probably valid for Fe-, Co-, Ni- and Mn-porphyrins, the second one - for metal free porphyrins, their complexes with Mg, Zn, Cd, Al, and Sn, chlorophylls and pheophytins. The rate constants of chemical quenching of singlet oxygen ( $K_{ox}$ ) were measured for the second group of compounds. The data suggest that chemical quenching is also a consequence of exciplex formation.

$K_q$  and  $K_{ox}$  values of carotenoids have been measured and investigated. The data suggest that formation of the exciplexes fully provides for the activity of vitamin A and retinal and for not less than 10% of activities of  $\beta$ -carotene and zeaxanthin.

The information obtained allow for a new interpretation of some concepts relevant to functioning of the quenchers in living cells. In particular, chlorophylls can be regarded not only as singlet oxygen photogenerators but also as photosynthetic membrane protectors.

1. Krasnovsky A.A., Jr. Biofizika USSR (1976) 21, 748; Photochem. Photobiol. (1979) 29, 29.

2. Krasnovsky A.A., Jr. Zh.Prikl.Spektrosk(1980) 32, 852; Chem.Phys.Lett. (1981) 81, 443; In: "Excited molecules. Kinetics of transformations (in Russian) Nauka, Leningrad (1982) 32; Photochem.Photobiol. (1982), 36,733; Krasnovsky A.A., Jr., Kagan V.E., Minin A.A. FEBS Lett. (1979) 108, 152; FEBS Lett. (1983) 155, 233; Krasnovsky A.A., Jr., Venedictov E.A., Chernenko O.M. Biofizika (1982) 27, 966; Krasnovsky A.A., Jr., Paramonova L.I. Biofizika (1983) 28, 725.



## SINGLET MOLECULAR OXYGEN IN PLANETARY ATMOSPHERES

E.J. Llewellyn and I.C. McDade  
Institute of Space and Atmospheric Studies  
University of Saskatchewan  
Saskatoon, Canada.

Airglow emissions from singlet molecular oxygen have been observed in the atmospheres of Earth, Mars and Venus although the emission intensities of the different systems are very different. Recent observations are reviewed and the vibrational development of the different emission systems discussed in some detail. A model which can describe this vibrational development is also presented.

The excitation mechanisms for the different emissions are considered and from a detailed examination of the volume emission profiles in the Earth's atmosphere it is shown that the emissions arising from the low lying singlet states are probably excited by energy transfer from some precursor. The required reaction rates are determined and it is suggested that the oxygen recombination reaction



leads to a preferential population of one electronic state. From a comparison between the Earth and Cytherean airglow it is proposed that the precursor, and the preferred state, is  $O_2(c \Sigma_u^-)$  and that the transfer rates are vibrational level dependent. An attempt to use a vibrational level independent scheme is also described and the resulting rate constants for the quenching of the precursor are determined. These nature of these values is to provide direct support for the laboratory measurements of Ogryzlo et al (1984).

E.A. Ogryzlo, Y.Q. Shen and P.T. Wassell, paper presented at COSMO 84.

t-Butylated hydroxytoluene sensitized photooxidation of the fatty acids: Inhibition by diazabicyclooctane and role of singlet oxygen.

M. K. Logani, P. D. Forbes and R. E. Davies, The Center for Photobiology, Skin and Cancer Hospital, Temple University Health Sciences Center, Philadelphia, PA 19140.

Environmental ultraviolet radiation, primarily of solar origin is generally conceded to be the primary etiologic agent for common forms of human skin cancer. In recent years, there has been accumulating evidence that antioxidants such as BHT, BHA, glutathione, etc. provide a considerable protection against photocarcinogenesis. The inhibitory action of antioxidants is believed to act through free radical quenching, thus preventing lipid peroxidation of cellular membranes. However, photooxidation of lipids in biomembranes is believed to occur through photosensitization via Type I or Type II mechanisms which are believed to be not affected by common antioxidants used to prevent 'dark oxidation'. The inhibitory effect of these antioxidants on photocarcinogenesis may thus be operating through some other mechanisms than by simple free radical quenching. In order to determine whether or not the phenolic antioxidants can prevent the photooxidation of lipids, we have studied the in vitro effect of some commonly used antioxidants on photooxidation of unsaturated fatty acids. Irradiation (wavelength > 320 nm) of methyl linoleate with unfiltered black light (10 x 20 watt) showed less than 1% consumption. A marked increase in the photooxidation (18%) of methyl linoleate was observed in presence of BHT. Similarly, consumption of methyl linolenate and arachidonate was much lower with black light alone (15.5 and 19.8% respectively) in comparison to when BHT was present (57.9 and 67.6% respectively). Conversely, BHA and tocopherol acetate did not show any significant effect on the photooxidation. Enhanced photooxidation by BHT was shown to involve  $^1O_2$  by characterization of the lipid hydroperoxides formed and by the effect of singlet oxygen quenchers. It was further shown that a diet supplemented with BHT and some other antioxidants enhanced photocarcinogenesis instead of reducing it. Whether this enhancement occurs through  $^1O_2$  is under further investigation.

Relaxation of the  $^1\Delta_g$  state in pure liquid oxygen  
and in liquid mixtures of  $^{16}\text{O}_2$  and  $^{18}\text{O}_2$ .

Max Maier

Naturwissenschaftliche Fakultät II - Physik,


Universität Regensburg

8400 Regensburg, Fed. Rep. Germany

We investigated the relaxation of the first excited electronic  $^1\Delta_g$  state and the vibrational levels of the electronic ground state  $^3\Sigma_g^-$  of oxygen in the liquid phase. The experiments were done in natural  $\text{O}_2$ ,  $^{18}\text{O}_2$ , and  $^{16}\text{O}_2$ - $^{18}\text{O}_2$  mixtures. The relaxation rate constants of the electronic states and the vibrational states were determined from time-resolved measurements of the fluorescence light and from time-delayed spontaneous anti-Stokes Raman scattering, respectively.

The following results were obtained:

- 1) The relaxation rate constants of the  $^1\Delta_g$  state in liquid natural  $\text{O}_2$  and  $^{18}\text{O}_2$  are  $2.2 \times 10^4 \text{ s}^{-1}$  and about  $100 \text{ s}^{-1}$ , respectively, at 77 K.<sup>1,2</sup> The relaxation rate is determined by intersystem crossing to higher vibrational levels of the electronic ground state



Calculations of the transition probability using the theory of Schwartz, Slawsky, and Herzfeld<sup>3</sup> (SSH) modified to include attractive forces<sup>4</sup> show, that the collisional deactivation of the  $^1\Delta_g$  state to the vibrational levels


$v=2$  and  $3$  of the electronic ground state should be dominant. The higher vibrational levels relax by rapid vibrational-vibrational energy exchange. In liquid natural  $O_2$  the energy accumulates in the  $^3\Sigma_g^-(v=1)$  state and is transferred slowly ( $\sim$  ms) to translational motion.

- ii) At high  $^1\Delta_g$  population additional relaxation channels cause a non-exponential decay of the  $^1\Delta_g$  state.<sup>2</sup> The energy pooling reaction to the second excited electronic  $^1\Sigma_g^+$  state of  $O_2$  is an important decay channel.
- iii) In  $^{16}O_2$ - $^{18}O_2$  mixtures there is a nonlinear dependence of the  $^1\Delta_g$  relaxation rate constant  $k$  on  $^{16}O_2$  concentration. For an explanation of the experimental results we discuss the following points: changes of the transition probability in the mixtures compared to the pure liquids, which are calculated from the SSH theory,<sup>3,4</sup> and the concentration dependence of the collision frequencies in the liquid mixtures.

1. E. Wild, H. Klingshirn, B. Faltermeier, and M. Maier, Chem. Phys. Letters 93, 490 (1982).
2. H. Klingshirn, B. Faltermeier, W. Hengl, and M. Maier, Chem. Phys. Letters 93, 485 (1982).
3. R.N. Schwartz, Z.I. Slawsky, and K.F. Herzfeld, J. Chem. Phys. 20, 1591 (1952).
4. H.K. Shin, J. Chem. Phys. 42, 59 (1965).

Structure and Kinetics of the Uncatalyzed Decomposition of Trialkyl Phosphite Ozonides. G.D. Mendenhall, R.F. Kessick, and M. Dobrzelewski, Department of Chemistry and Chemical Engineering, Michigan Technological University, Houghton, MI 49931.

Activation parameters for uncatalyzed loss of oxygen from phosphite ozonide solutions (equation (1)) were measured for a number of different



phosphites in several solvents. Triphenyl phosphite ozonide (TPPO) in toluene, chlorobenzene, and acetonitrile showed  $E_a$  (kcal/mole) = 19.4-20.1, and  $\log(A/\text{sec}) = 13.5-13.6$ , whereas in dichloromethane the rate constants were not very reproducible and depended on the precautions followed to exclude moisture from the ozonide preparations. Activation parameters for TPPO in dichloromethane reported in the literature, or derived from reported rate constants vary from  $\log(A/\text{sec}) = 8.1-11.9$  and  $E_a$  (kcal/mole) = 14.4-17.2. Other phosphite ozonides show activation parameters which range approximately between the extremes given above for TPPO, even in solvents other than dichloromethane. We suggest that the decompositions with relatively high  $E_a$  and  $\log A$  proceed through a singlet biradical intermediate, and those with low and variable activation parameters involve a zwitterion or an induced decomposition. The greater stability of cyclic and bicyclic ozonides is suggested to result in part from O-O bond strengthening in the cyclic cases, analogous to O-H bond strengthening in certain bicyclic hydroxylamines. An earlier theory involving O-O bond cleavage in the ozonide and pseudorotation of the resulting open-chain intermediate is discussed. A molecular weight determination and the  $31\text{P-NMR}$  spectrum of trimethylol-phosphite ozonide are consistent with a monomeric, non-equilibrating structure.

Singlet Oxygen Oxidation of Substituted Thiobenzamides. Robert W. Murray  
and Sudhir K. Agarwal, Department of Chemistry, University of Missouri-  
St. Louis, St. Louis, Missouri 63121.

Thiobenzamides are believed to require metabolic activation in order to express their observed hepatotoxicity. Hanzlik and coworkers have shown that relative toxicity, using several indices of toxicity in vivo, can be correlated with thiobenzamide substituent Hammett  $\sigma$  values in a manner which is consistent with an electrophilic biooxidation process. They have also found a similar  $\rho\sigma$  correlation when using substitute thiobenzamides and  $H_2O_2$  as the oxidant. During the course of our studies on the use of carbonyl oxides as models for the flavin-assisted mono-oxygenases we have had reason to study the reactions of substituted thiobenzamides with singlet oxygen. The relative rates of these reactions have been found to fit a Hammett  $\rho\sigma$  correlation with  $\rho = -0.39$  indicative of an electrophilic oxidation process.

Thursday

POSSIBLE ROLE OF SINGLET OXYGEN IN THE CONTROL OF THE PHOTOTACTIC REACTION  
SIGN OF THE CYANOBACTERIUM ANABAENA VARIABILIS.

Wilhelm Nultsch and Hartwig Schuchart, Department of Botany, University of Marburg, FRG.

In earlier papers (Nultsch et al., Arch.Microbiol. 134, 33-37, 1983; Nultsch and Wenderoth, Z.Pflanzenphysiol. 111, 1-7, 1983) it has been shown that the phototactic reaction sign of the cyanobacterium Anabaena variabilis at high fluence rates is reversed from negative (movement away from the light source) to positive (movement toward the light source) by addition of  $10^{-3}$  M sodium azide. Moreover, photobleaching is also prevented by  $10^{-3}$  M azide (Nultsch et al., Z.Pflanzenphysiol. 111, 327-332, 1983). Based on these findings the hypothesis has been proposed that the phototactic reaction sign reversal generator of Anabaena is controlled by the level of an active oxygen species, probably singlet oxygen, which is quenched by azide.

In order to confirm this hypothesis, the effects of other quenching agents have been studied.  $5 \cdot 10^{-3}$  M DABCO (= diazabicyclo-2,2,2-octane) reverses the reaction sign from negative to slightly positive, but is cytotoxic at higher concentrations. L-Histidin inhibits negative phototaxis at concentrations  $< 5 \cdot 10^{-4}$  M, whereas positive phototaxis is not influenced. On the contrary  $2 \cdot 10^{-2}$  KJ inhibits both positive and negative phototaxis. This could be due to a quenching of the photoreceptor triplet state.

Experiments with phosphate buffer of different pH demonstrate that the transition point from positive to negative phototaxis is shifted from  $2.7 \text{ Wm}^{-2}$  at pH 7 to  $10 \text{ Wm}^{-2}$  at pH 6, and that at pH 5.5 only positive responses occur even at  $50 \text{ Wm}^{-2}$ . At lower pH values movement ceases because of the decrease of the proton motive force. If imidazole buffer (pH 6,0) is used, negative phototaxis is suppressed the more the higher the imidazole concentration is. Above  $2.5 \cdot 10^{-4}$  M positive reactions occur at  $13,5 \text{ Wm}^{-2}$ .


These results as well as some experiments being in progress support the hypothesis that the singlet oxygen level regulates the phototactic reaction sign, probably via a reaction sign reversal generator, this way enabling the Anabaena filaments to escape dangerous light conditions by negative phototaxis.

E.A. Ogryzlo, Y.Q. Shen and P.T. Wassell


Department of Chemistry,  
University of British Columbia  
Vancouver, Canada

Earlier laboratory studies of the yield of  $O_2(b^1\Sigma_g^+)$  in the recombination of oxygen atoms in the gas phase resulted in values which were too small to account for the observed emission from  $O_2(b)$  in the atmospheric night airglow. We have reinvestigated this process in a fast flow system using the  $N+NO$  reaction as a source of oxygen atoms. The rate of formation of  $O_2(b)$  along the observation tube was monitored by following the 762 nm emission from this state.

In the absence of any added  $O_2$ , the production of  $O_2(b)$  in the stream of oxygen atoms was found to occur even more slowly than indicated by earlier work. When  $O_2$  was added to the system a dramatic increase in the  $O_2(b)$  production rate was observed. We have measured the kinetic dependencies of the emission intensity when a steady state concentration of  $O_2(b)$  is reached. These results indicate that the process cannot be described by the simple termolecular reaction:



The observations are, however, consistent with the mechanism described by reactions 1-4, in which recombination initially occurs into a precursor ( $O_2^*$ ) which is quenched by oxygen atoms and yields  $O_2(b)$  only on collision with  $O_2$ .



When the  $O_2$  concentration is large reaction 1 is rate controlling and  $k_1$  can be calculated from a measurement of the steady state  $O_2(b)$  concentration and an independent measurement of  $k_4$ . The value which we obtain for  $k_1$  ( $1.6(\pm 0.2) \times 10^{-34} \text{ cm}^6 \text{ molec}^{-2} \text{ s}^{-1}$ ) is confirmed by separate measurements of the rise time for this state.

The identity of  $O_2^*$  remains uncertain. Our results indicate that it cannot be identified with the lower vibrational levels of  $O_2(A)$ ,  $O_2(A')$  or  $O_2(c)$  since the ratio  $k_2/k_3$  for  $O_2^*$  is  $3.6 \pm 0.7$  which is much too small for these states.



Optical Determination of the Rates of Formation and Decay of  $O_2(^1\Delta_g)$   
in  $H_2O$ ,  $D_2O$  and Other Solvents

J. G. Parker and W. D. Stanbro  
The Johns Hopkins University  
Applied Physics Laboratory  
Laurel, Maryland 20707

Abstract

Recent experimental investigation, using dissolved photosensitizers subjected to pulsed laser radiation, has permitted accurate determination of the quenching rates of  $O_2(^1\Delta_g)$  in a wide variety of solvents. Of particular importance in this application is the use of remote optical sensing to monitor the temporal behavior of the resultant  $O_2(^1\Delta_g)$  concentration produced.<sup>1-3</sup> Such optical sensing involves detection of the relatively weak emission accompanying the collisionally induced radiative transition to the  $^3\Sigma_g^-$  ground electronic state. One distinct advantage inherent in this method, in addition to the fact that it obviates problems associated with alternatives such as chemical acceptors, is that a positive identification of the emitting species as  $O_2(^1\Delta_g)$  is possible through determination of the emission spectrum.

Another important advantage of remote optical monitoring is that  $O_2(^1\Delta_g)$  concentrations may be determined in a heterogeneous medium,<sup>4</sup> one of the most important applications in this regard being the interface separating a lipid and aqueous phase. It is clear that rather significant  $O_2(^1\Delta_g)$  concentration gradients at this interface must occur due to the fact that in general lifetimes in  $H_2O$  are approximately an order of magnitude shorter than in a hydrocarbon environment. Recent studies<sup>5</sup> have shown that the quenching rate for a given solvent depends strongly on the presence of bound H atoms and in particular on the exact nature of the binding, i.e., C-H or O-H type.

The relatively long decay times in hydrocarbon solvents provide a particularly favorable case for optical monitoring. Thus, for example, the lifetime of  $O_2(^1\Delta_g)$  in benzene has been established as lying in the range 30-35  $\mu\text{sec}$ . Addition of OH groups causes a systematic lowering of lifetimes, e.g., 15  $\mu\text{sec}$  for ethanol and 48  $\mu\text{sec}$  for ethylene glycol.<sup>4</sup> Earlier measurements of the lifetimes in  $H_2O$  have ranged from 2-7  $\mu\text{sec}$ . Factors tending to complicate lifetime measurements in this important case are twofold: (1) relatively long (2-3  $\mu\text{sec}$ ) times required for transfer of energy from the sensitizer triplet, and (2) interference due to the presence of an infrared component of the sensitizer fluorescence. Separation of the latter from the  $O_2(^1\Delta_g)$  emission places a premium on fast detector response time. Earlier measurements indicated that germanium photodiodes used as detectors were inadequate in this regard. Recent measurements, however, using an InGaAs device have yielded much more satisfactory results.<sup>4</sup>

The problem of determination of the lifetime of  $O_2(^1\Delta_g)$  in  $H_2O$  in addition to requiring detectors capable of rapid response also requires use of a sensitizer capable of providing a high  $O_2(^1\Delta_g)$  yield and exhibiting high resistance to subsequent oxidation. Evaluation of a series of water soluble synthetic porphyrins led to the selection of tetra(4-N-methylpyridil) porphyrin as being the optimum sensitizer. This sensitizer, soluble in  $H_2O$  over a wide range of

Wednesday

pH, proved extremely resistant to degradation even when incident laser pulse energies of 100 millijoules were used, ten times the level used normally. Possibly this inertness is due to an absence of photoionization processes.<sup>6</sup>

A detailed series of measurements was carried in H<sub>2</sub>O using a T4MPP sensitizer for air-saturated, oxygen-enriched and oxygen-depleted samples, for low, medium and high pulse energies. No systematic dependence of decay time on laser pulse energy was observed; however, a definite shortening of the formation time in air-saturated solutions was observed with increasing laser power. Due to the fact that the triplet transfer times and O<sub>2</sub>(<sup>1</sup>Δ<sub>g</sub>) decay times were very close to the same value for the case of air saturation, it was impossible to carry out an accurate determination of the respective time constants. Thus, determination of O<sub>2</sub>(<sup>1</sup>Δ<sub>g</sub>) decay times necessitated the use of oxygen-enriched samples with correspondingly reduced triplet energy transfer times. Statistical averaging of the decay times for 14 independent runs gave the value  $\tau_D = 3.21 \pm .12 \mu\text{sec}$ . In all these runs the sensitizer concentration was fixed at  $1 \times 10^{-4} \text{M}$ . Similar runs with the concentration reduced to  $5 \times 10^{-5} \text{M}$  produced results in agreement with this value. Using this value for the O<sub>2</sub>(<sup>1</sup>Δ<sub>g</sub>) decay time, triplet transfer times were extracted from the formation times characterizing the air-saturation cases. Analysis of the data gives  $\tau_T = 2.43 \pm .18 \mu\text{sec}$ .

Similar runs were carried out using the same sensitizer in D<sub>2</sub>O. In this case the O<sub>2</sub>(<sup>1</sup>Δ<sub>g</sub>) decay time is greatly lengthened while the triplet transfer time is essentially unchanged thus permitting the use of air saturated solutions. Analysis of the data yielded  $\tau_D = 68 \mu\text{sec}$  and  $\tau_T = 2.1 \mu\text{sec}$ . Thus solvent deuteration gives rise to an increase in the O<sub>2</sub>(<sup>1</sup>Δ<sub>g</sub>) decay time of more than an order of magnitude. It is interesting to point out that earlier measurements using another water-soluble synthetic porphyrin tetra(4-sulfonato-phenyl)porphine gave for D<sub>2</sub>O an initial value of 55  $\mu\text{sec}$ , however, this value steadily decreased with increasing length of exposure to the laser radiation. This sensitizer degradation is possibly due to its known ability to photoionize<sup>6</sup> coupled with the relatively long O<sub>2</sub>(<sup>1</sup>Δ<sub>g</sub>) lifetime. No such effects were observed using the T4MPP sensitizer.

Other solvents investigated were: ethylene glycol, characterized by increased viscosity and correspondingly lengthened triplet transfer times; 1-octanol with similar behavior, however with a significantly longer O<sub>2</sub>(<sup>1</sup>Δ<sub>g</sub>) decay time; glacial acetic acid and formamide.

#### References

1. J. R. Hurst, J. D. McDonald and G. B. Schuster, J. Am. Chem. Soc. 104, pp. 2065-2067 (1982).
2. J. G. Parker and W. D. Stanbro, J. Am. Chem. Soc. 104, pp. 2067-2069 (1982).
3. P. R. Ogilby and C. S. Foote, J. Am. Chem. Soc. 104, pp. 2069-2071 (1982).
4. J. G. Parker and W. D. Stanbro, Proceedings of the Clayton Foundation Symposium on Porphyrin Localization and Treatment of Tumors, (D. R. Doiron, Ed.) A. R. Liss, Inc. (1983).
5. J. R. Hurst and G. B. Schuster, J. Am. Chem. Soc. 105, pp. 5756-5760 (1983).
6. R. Bonnet, R. J. Ridge, E. J. Land, R. S. Sinclair, D. Tait and T. G. Truscott, J. Chem. Soc., Faraday Trans. I, 78, pp. 127-136 (1982).

Friday

Optical Determination of  $O_2(^1\Delta_g)$  Quenching Rates and Relative  
Emission Intensities in High Pressure Oxygen Gas Using  
Pulsed Laser Excitation at  $1.064 \mu$

J. G. Parker  
The Johns Hopkins University  
Applied Physics Laboratory  
Laurel, Maryland 20707

Abstract

Singlet molecular oxygen in the  $v' = 1$  vibrational level of the  $^1\Delta_g$  electronic state was generated in high pressure oxygen gas by means of pulsed laser radiation at a wavelength of  $1.064 \mu$ . The resulting  $O_2(^1\Delta_g)$  population was monitored optically, using a remotely positioned germanium photodiode in combination with appropriate filtering to record the subsequent infrared emission. Pressures ranged from approximately 40 to 95 atm. Measurements were carried out both in the gas phase and also with the oxygen dissolved in a freon solvent.

The observed time dependence of this infrared emission in the gas phase is well described by an exponential with an average time constant  $p_o\tau = .0221$  sec-atm. Runs were also made in which a sizable helium addition was used to investigate possible effects due to an increased vibrational quenching rate on the overall  $O_2(^1\Delta_g)$  deactivation process. A slight increase in the time constant to the value .0250 sec-atm was found, comparing favorably with the value .0244 sec-atm obtained from earlier measurements on a similar mixture using a photoacoustic technique.<sup>1</sup> Lifetimes obtained by others using infrared monitoring techniques, in which significantly different methods of  $O_2(^1\Delta_g)$  generation were used, lie in the range from .0179 to .0259 sec-atm, encompassing the above values.

A series of runs was devoted to a determination of the dependence of emission intensity on oxygen pressure, with peak intensity being recorded as the pressure was increased sequentially from an initial value of 47.9 atm to a final value of 80.9 atm. A plot of the logarithm of relative emission intensity  $I_R$  versus logarithm of pressure  $p_o$  indicated that  $I_R = Kp_o^3$ . It is well known that the optical absorption at  $1.064 \mu$  leading to the formation of  $O_2(^1\Delta_g)$  varies quadratically with pressure and the presence of the additional factor thus implies that the emission rate must depend linearly on  $p_o$  in accord with experimental results obtained at lower pressure. The emission process at these pressures is therefore collisionally induced. Calculation, using the empirical dependence of radiative lifetime on gas pressure established by Badger et al.<sup>2</sup> at lower pressures, indicates that in the pressure interval from 50 to 100 atm, this lifetime should range from 10 to 20 seconds.

Recent data indicates  $O_2(^1\Delta_g)$  radiative lifetimes in various solvents to actually be independent of the particular solvent and to be equal to a value of approximately 4 seconds.<sup>3</sup> The suggestion is therefore that for solvents exhibiting high oxygen solubility, such as the freons, sequential pressurization to levels approaching 100 atm from lesser values ought to span an important range, at the lower end of which emission due to  $O_2(^1\Delta_g)$ -solvent collisions is dominant, to the upper end where  $O_2(^1\Delta_g)$ -oxygen collisions

control. At lower pressures, therefore, a quadratic dependence of emission intensity on pressure is expected, while at higher pressures transition to a cubic dependence should be observed.

Thus a series of runs was carried out in which the cylindrical observation section of a high pressure cell was filled with Freon-113 (1,1,2-trichlorotrifluoroethane) and the overlying volume carefully pressurized with oxygen. The laser was positioned so that its beam axis coincided with that of the observation section of the cell. With the laser incident pulse energy fixed at 40 millijoules,  $O_2(^1\Delta_g)$  decay curves were obtained at time intervals of one hour subsequent to pressurization. Corresponding decay times and amplitudes were obtained from each run. These times were observed to decrease steadily as oxygen diffused to the center of the cell, with an increase in the amplitude factor.

Equilibrium was attained after a period of eight hours. The dependence of the reciprocal equilibrium time constants on oxygen pressure was linear, with a slope consistent with the gas phase data and an intercept of 1.30 (millisec)<sup>-1</sup> corresponding to a time constant of 770  $\mu$ sec, i.e., the deactivation time for  $O_2(^1\Delta_g)$  infinitely dilute in the Freon-113 solvent. This value is substantially smaller than that obtained for this solvent under air-saturated conditions.

Dissolved oxygen levels were determined as a function of time from the sequential values of decay times obtained during the approach to equilibrium. Using a simplified model of the cell geometry, axial oxygen levels were calculated as a function of time from a one-dimensional diffusion equation. Comparison of results of these calculations with the experimental data yielded a diffusion coefficient  $D = 5.8(-5)$  cm<sup>2</sup>/sec.

The dependence of emission intensity on oxygen pressure exhibited a significant departure from a cubic dependence at the lowest pressures indicating the dominance of  $O_2(^1\Delta_g)$ -solvent collisions in this region. However, the onset of the  $O_2(^1\Delta_g)$ -oxygen collision controlled regime occurs at lower values of pressure than one would predict from lifetime calculations based on Ref. 2.

#### REFERENCES

1. J. G. Parker, J. Chem. Phys. 67, pp. 5352-5361 (1977).
2. R. M. Badger, A. C. Wright and R. F. Whitlock, J. Chem. Phys. 43, pp. 4345-4350 (1965).
3. A. A. Krasnovsky, Jr., Chem. Phys. Ltrs. 81, -p. 443-445 (1981).
4. B. Stevens and K. L. Marsh, J. Phys. Chem. 86, pp. 4473-4476 (1982).

## Reactive Oxygen Species in the Biological Effects of Solar Ultraviolet Radiation.

Meyrick J. Peak,\* Jennifer G. Peak\* and Christopher S. Foote.<sup>†</sup>

\*Division of Biological and Medical Research, Argonne National Laboratory, Argonne, IL 60439, and <sup>†</sup>Department of Chemistry, University of California, Los Angeles, CA 90024.

Ultraviolet action spectra for the killing of *Escherichia coli* by ultraviolet (UV) and visible radiations in the presence and absence of ground state oxygen were prepared. No oxygen enhancement ratios (OERs) were obtained at wavelengths shorter than 320 nm. Above this wavelength, we observed OER values that increased with increasing wavelength up to a value of about eight at 405 nm (1). Our hypothesis of the mechanism whereby molecular oxygen participates in the lethal effects of these longer wavelengths is that the radiation reacts with an endogenous photosensitizer, which in turn excites ground state oxygen into one or more reactive oxygen species, which then kill the cell. It is generally believed that it is damage to DNA that is the critical cause of this cell killing (2). We have identified one of the cellular sensitizers involved in these lethal events in *E. coli* as 4-thiouridine ( $s^4U$ ), a rare base found in tRNA. This nucleoside is responsible for 20 to 50% of the lethal events caused by 334 nm UV radiation (3).

In order to avoid problems with permeability barriers in the cells, we used purified genetically active DNA from *B. subtilis* in our studies. We measured its genetic activity and molecular weight after treatment with near-UV radiation and a variety of added modifying compounds (quenchers and sensitizers). The following results were obtained. Experiments in which the purified DNA was irradiated anoxically by monochromatic near-UV radiation ( $> 320$  nm) showed that oxygen plays a role in the loss of DNA genetic activity and also in the induction of DNA single-strand breaks (SSBs). Any of the quenchers--histidine, diazobicyclo[2.2.2]octane (DABCO), aminoethylisothiouonium bromide hydrobromide (AET), benzoate, azide, and glycerol--added during irradiation were shown to protect the DNA against near-UV radiation damage (see 2 for review). Replacement of  $H_2O$  with  $D_2O$  during the irradiation doubled the rate of induction of SSBs in the DNA. Our UV action spectra for the modifying effects (histidine, AET, DABCO, glycerol) all show maximum protection near 350 nm. At 300 nm and below, no protective effect is shown. In other experiments, a number of putative sensitizers were added to the DNA during irradiation. These compounds were the free nucleosides  $s^4U$  and 5-methylaminomethyl-2-thiouridine, the free base 2-thiouracil ( $s^2Ura$ ), the nucleotide coenzyme riboflavine, and the oxidized and reduced forms of nicotinamide adenine dinucleotide. All these compounds increased the induction of SSBs in DNA by monochromatic 334 nm UV radiation (4). As  $s^2Ura$  was the most effective photosensitizer for induction of the SSBs, it was selected for further study. Maximal enhancement by  $s^2Ura$  of SSB induction was shown to occur when  $s^2Ura$  was present at a concentration of approximately 0.001 M. Anoxia, histidine, DABCO, benzoate and glycerol when present separately during irradiation in the presence of 0.001 M  $s^2Ura$  were all shown to decreased the number of SSBs caused by  $s^2Ura$  and 334 nm UV radiation. Studies were performed of the concentration dependence of the protection by DABCO and by benzoate against the SSBs caused by 334 nm radiation in the

presence of 0.001 M  $s^2$ Ura. The sigmoidal shapes of these curves showed maximal protection by DABCO and benzoate at a concentration of 0.1 M against SSBs caused by 334 nm radiation in the presence of  $s^2$ Ura and no protection below a concentration of 0.001 M. When  $H_2O$  was replaced by  $D_2O$  during irradiation of the DNA by 334 nm radiation in the presence of  $s^2$ Ura, the rate of induction of SSBs was about doubled. Finally, superoxide dismutase reduced by about 20% the rate of induction of SSBs in the DNA by 334 nm radiation in the presence of 0.001 M  $s^2$ Ura.

(This work is supported by the U.S. Department of Energy under contract No. W-31-109-ENG-38.)

#### References

1. Peak, J. G., Peak, M. J., and Tuveson, R. W. (1983) Photochem. Photobiol., in press.
2. Peak, M. J. and Peak, J. G. (1982) NATO Advanced Res. Inst. Monograph "Biological Effects of Solar UV Radiation." Plenum Press, New York, pp. 325-336.
3. Peak, M. J., Peak, J. G., and Nerad, L. (1983) Photochem. Photobiol. 37, 169-172.
4. Peak, J. G., Peak, M. J., and MacCoss, M. (1983) Photochem. Photobiol., in press.

SYNERGESTIC AND ANTAGONISTIC EFFECTS OF QUENCHING  
SINGLET OXYGEN IN A POLYMER MATRIX

B.Rånby and J.F.Rabek

Department of Polymer Technology, The Royal  
Institute of Technology, Stockholm, Sweden

A deterioration of many polymeric materials (rubbers, plastics, fibers and biopolymers) occurs under sunlight irradiation in photooxidation processes (light ageing processes) in which a singlet oxygen  $^1\text{O}_2$  may play a considerable role <sup>1</sup>. The most susceptible polymers towards  $^1\text{O}_2$  oxidation are polydienes <sup>2-4</sup>. The reaction of  $^1\text{O}_2$  with polybutadiene (PB) leads to the formation of allylic hydroperoxide according to the ene-type mechanism.

Various compounds have been reported to quench  $^1\text{O}_2$  in the solution or polymer matrix. These compounds include carotenoids <sup>5</sup>, hindered piperidines <sup>6</sup>, phenols <sup>7</sup>, etc. However, all of these compounds are either inefficient quenchers or are themselves unstable and subject to photooxidation.

In this paper we present results of the synergistic and antagonistic effects of quenching of  $^1\text{O}_2$  by the mixture of hindered phenols and hindered piperidines in a PB matrixes and in solution.

It is generally accepted that the presence of carbonyl groups (internal and/or external impurities) in a polymer matrix can be responsible for the generation of  $^1\text{O}_2$  (energy transfer mechanism). Considering a fact that triplet-excited benzophenone can produce  $^1\text{O}_2$  <sup>8</sup> we have studied its reaction with PB. Hindered phenols beside their reactions with  $^1\text{O}_2$  may also react with triplet-benzophenone to yield fuchsone derivatives. Addition of hindered phenols decreases sensitized photooxidation of PB. A mechanism of these reactions is discussed in a detail.

References

1. J.F.Rabek, in "Singlet O<sub>2</sub>" (ed.A.A.Frimer),CRC (in press).
2. J.F.Rabek and B.Rånby, Photochem.Photobiol.,28,557 (1978).
3. J.F.Rabek ,J.Lucki and B.Rånby, Europ.Polym.J.,15,1089 (1979).
4. M.A.Golub,NASA Technical Memorandum,No.78604,1979.
5. J.F.Rabek and D.Lala, J.Polym.Sci.,B18,427 (1980).
6. Y.Y.Yang,J.Lucki,J.F.Rabek and B.Rånby, Polym.Photochem.,3,  
97 (1983).
7. J.F.Rabek,B.Rånby and J.Arct, Polym.Degrad.Stabil.,5.65 (1983).
8. A.M.Trozzolo in "Oxidation of Organic Compounds",Vol.III,p.167,  
Adv.Chem.Ser.,No.77,ACS,Washington,D.C.1968.



Michael A. J. Rodgers- CFKR, University of Texas, Austin, TX 78712

"The Relaxation of  $O_2(^1\Delta_g)$  in Condensed Systems"

This paper will serve largely as a brief introductory statement for the session on "Relaxation and Quenching." The major focus will be on the non-radiative decay of  $O_2(^1\Delta_g)$  in liquid systems as determined through measurements of the radiative process  $^3\Sigma_g^- \leftarrow ^1\Delta_g$  that result in luminescence at 1.27  $\mu\text{m}$ . Time resolution of this luminescence to the submicrosecond regime has been achieved by the use of high intensity laser pulses, fast photodetectors and high bandwidth amplifiers. This has led to rapid, straightforward and precise measurement of  $O_2(^1\Delta_g)$  lifetime values ( $\tau_\Delta$ ) in a variety of solvent systems. Recent work has shown that bimolecular quenching constants for solvent quenching exhibit regular variations throughout homologous series of n-alkyl compounds and additivity parameters can be evaluated. This leads to the possibility that  $\tau_\Delta$  values can be predicted from a consideration of molecular structure. Other data will be presented on the non-radiative decay of  $O_2(^1\Delta_g)$  in micro-heterogeneous liquid systems and water-swollen polymers.

ON THE USE OF WATER-SOLUBLE  $^1\text{O}_2$  SOURCE FOR MECHANISTIC STUDY AND FOR  
OXYGENATION OF BIOLOGICAL SYSTEMS


Isao Saito, Kenzo Inoue, Hiroshi Sugiyama, and Teruo Matsuura

Department of Synthetic Chemistry, Faculty of Engineering, Kyoto University  
Kyoto 606, Japan

We observed the formation of  $\text{O}_2^-$  from the reaction of electron-rich substrates, e.g., N,N-dimethylanilines and  $^1\text{O}_2$  generated from a water-soluble endoperoxide 1 in aqueous media.<sup>1,2</sup> We suggested that the substrates with oxidation potentials less than 0.5 V vs. SCE are capable of undergoing one electron-transfer to  $^1\text{O}_2$  to generate  $\text{O}_2^-$  in polar aqueous solvents.<sup>2</sup> In order to know the generality of  $\text{O}_2^-$  formation, we have examined the reaction of thioanisoles with  $^1\text{O}_2$  in aqueous solvents. p-Hydroxythioanisole ( $E_{1/2} = 0.48$  V vs. SCE) was readily oxidized to the corresponding sulfoxide at 35 °C in the presence of 1 in aqueous methanol. In this reaction the formation of  $\text{O}_2^-$  was confirmed by the assay utilizing a combination of NBT and SOD. Addition of increasing amounts of NBT reduced the yield of sulfoxide with the enhancement of NBT reduction. The result suggests that  $\text{O}_2^-$  is directly involved in the sulfoxidation.

Other example for the use of the water-soluble  $^1\text{O}_2$  source is the oxidative cleavage of DNA.  $^1\text{O}_2$  is known to selectively oxidize guanine base among nucleic acid bases. We have synthesized a new intercalator (2) possessing  $^1\text{O}_2$  source 1 at the side chain. Incubation of a 5'-[ $^{32}\text{P}$ ]-end-labeled DNA fragment with 2 at 35 °C in phosphate buffer (pH 7) followed by piperidine treatment gave rise to the efficient cleavage of the DNA chain selectively at guanine residues.

Usefulness of polymer-bound  $^1\text{O}_2$  source in oxygenation of organic substrates will also be discussed.



1. I. Saito, T. Matsuura and K. Inoue J. Am. Chem. Soc. 103, 188 (1982).
2. I. Saito, T. Matsuura and K. Inoue ibid., 105, 3200 (1983).


## COSENSITIZED ELECTRON-TRANSFER PHOTOOXYGENATION

A. Paul Schaap,\* Shahabuddin Siddiqui,  
Girija Prasad, and Luigi Lopez,

Department of Chemistry, Wayne State University

Detroit, Michigan 48202

9,10-Dicyanoanthracene (DCA) and biphenyl (BP) cosensitize the electron-transfer photooxygenation of epoxides in oxygen-saturated acetonitrile to form ozonides. These epoxides do not measurably quench the fluorescence of DCA and are, therefore, relatively unreactive under standard photooxygenation conditions with only DCA as sensitizer. However, a dramatic increase in the rate of reaction is observed upon the addition of BP as a non-light-absorbing cosensitizer. An investigation of the stereochemistry of the reaction with *cis*- and *trans*-2,3-diphenyloxiranes has shown that both epoxides are converted exclusively to the *cis*-ozonide. This striking stereochemical result is consistent with a proposed mechanism involving singlet oxygen addition to an intermediate carbonyl ylide.



DCA-sensitized photooxygenation of aziridines provides the corresponding 1,2,4-dioxazolidines. The addition of biphenyl as a cosensitizer is not required in this case as the more easily oxidized aziridines efficiently quench the fluorescence of DCA. The stereochemistry of the product is a function of R. The *cis* isomer is virtually the exclusive product when  $\text{R}=\text{H}$ . However, the *trans* isomer is the major product when  $\text{R}=\text{PhCH}_2$ . A mechanism involving the addition of singlet oxygen to intermediate azomethine ylides is proposed.


NUCLEOPHILIC OXYGEN-ATOM TRANSFER FROM A  
PEREPOXIDE INTERMEDIATE TO SULFOXIDES

A. Paul Schaap,\* Scott G. Recher,  
Gary R. Faler, and Steven R. Villasenor

Department of Chemistry, Wayne State University,  
Detroit, Michigan 48202

Singlet oxygen reacts with adamantylideneadamantane to form an unusually stable 1,2-dioxetane. We have recently obtained evidence for a trappable intermediate in this 1,2-cycloaddition reaction of  $^1\text{O}_2$ . This species, which we propose is the perepoxi intermediate shown below, can be quantitatively trapped by nucleophilic oxygen-atom transfer to sulfoxides to yield sulfones and the epoxide. The ratio of epoxide to dioxetane is found to be a function of the concentration of the sulfoxide.

Additional evidence for the proposed mechanism is provided by a study of substituent effects on the trapping reaction. A series of competition experiments with substituted-phenyl methyl sulfoxides gave a Hammett plot with a  $\rho$  value of +0.52. These results are consistent with the proposed nucleophilic character of the intermediate. Solvent effects on this reaction have also been investigated.




SELF-SENSITIZED PHOTOOXIDATION OF AROMATIC COMPOUNDS AND  
PHOTOCYCLOREVERSION OF ENDOPEROXIDES.  
APPLICATIONS IN CHEMICAL ACTINOMETRY.

R. Schmidt and H.-D. Brauer


Institute for Physical and Theoretical Chemistry,  
University of Frankfurt, Robert-Mayer-Str. 11,  
D 6000 Frankfurt/Main, F.R.G.

The self-sensitized photooxidation of aromatic compounds (HC) leading with quantum yield  $Q_2$  to the formation of thermally stable endoperoxides (HCPO) is a photoreversible process. Via adiabatic photocycloreversion the endoperoxides can be split into the parent aromatic compound and singlet oxygen during short wavelength irradiation<sup>1,2</sup>. Therefore, the HCPO/HC + O<sub>2</sub> systems are in principle photochromic. However, irreversible rearrangement reactions (quantum yield:  $Q_{dec}$ ) of the endoperoxides generally compete with cycloreversion, thus reducing the number of possible photochromic cycles<sup>3</sup>.



Due to the large shift in wavelength of absorption, which accompanies the photooxidation and the photocycloreversion, the HCPO/HC + O<sub>2</sub> systems are in principle well suited as convenient chemical actinometers, in which the absorbed radiation is determined spectrophotometrically. However, a photochemical system has to meet a number of additional important requirements in order to be a useful chemical actinometer. These are: exact knowledge of the relevant photochemical reactions; reproducibility; almost wavelength independent actinometric factors; high sensitivity; high accuracy; and simplicity in the evaluation of the data.


It will be demonstrated that the self-sensitized photooxidation of mesodiphenylhelianthrene (MDH) fulfils these requirements. Above all MDH is well suited as actinometric material



because the quantum yield of photooxidation  $Q_2$  depends neither on the irradiation wavelength nor on the concentrations of MDH and O<sub>2</sub> in the ranges used for actinometry.

- 3 -

The highly reversible photochromic system consisting of hetero-coerdianthrone (HCD) and its endoperoxide (HCDPO) is excellently suitable for reusable actinometry in the uv-region. During



each actinometric measurement the original colourless solution turns red due to the formation of HCD. Subsequently, the solution can be decolourized completely by using a cut off filter and sun light, thus being ready again for the next experiment. In this way it is possible to do 100 actinometric measurements with the same solution without loss in accuracy.

<sup>1</sup>R. Schmidt, W. Drews and H.-D. Brauer, J. Am. Chem. Soc., 102, 2971 (1980)

<sup>2</sup>R. Schmidt, J. Photochem., 23, 379 (1983)

<sup>3</sup>R. Schmidt, W. Drews and H.-D. Brauer, J. Photochem., 18, 365 (1982)

POSSIBLE FORMATION OF SINGLET OXYGEN FROM  
VIBRATIONALLY EXCITED WATER

Ajit Singh, Grant W. Koroll and Stephen A. Antonsen

Medical Biophysics Branch  
Atomic Energy of Canada Limited Research Company  
Whiteshell Nuclear Research Establishment  
Pinawa, Manitoba, Canada ROE 1L0

Solubility of oxygen in  $H_2O$  and  $D_2O$  is similar and is lower than that in Freon-113 by a factor of about 40. In a 5 cm cell in solutions in Freon-113 ( $\sim 13$  MPa), absorption bands of oxygen are clearly observed in the wavelength range 370-1300 nm. However, in solution in water at that pressure only the absorption band at 760 nm is observed.

In the literature solute decomposition has been attributed to singlet oxygen for solutions of  $O_2$  in  $D_2O$  ( $\sim 15$  MPa) on irradiation at 1056 nm, with a Nd-Yag laser. However, under these conditions absorption of light by  $D_2O$  would be about two orders of magnitude greater than that by oxygen. This raises the question, how is singlet oxygen formed in solutions of  $O_2$  in  $D_2O$  ( $\sim 15$  MPa)?

Solutions of 1,3-diphenylisobenzofuran (DPBF,  $\sim 1 \times 10^{-5}$  mol  $dm^{-3}$ ) in  $H_2O$  and  $D_2O$  (containing 3% acetonitrile), under 13 MPa oxygen, were partially bleached on exposure to repeated pulses from a Nd-Yag laser. The bleaching was greater in the solution in  $H_2O$ . Similar bleaching was also seen when DPBF solutions (open to air) were exposed to light at 990 nm from a 1600 W Xe-arc lamp. Decomposition of solutions of dimethylfuran ( $1 \times 10^{-3}$  mol  $dm^{-3}$ ) in  $H_2O$  and  $D_2O$  (open to air) was also seen on exposure to light,  $\lambda > 850$  nm, from a Xe-arc lamp, but the decomposition appeared to be greater in the solutions in  $D_2O$ .

The results suggest that one or both of the following mechanisms contribute to solute decomposition in aqueous solutions irradiated at  $\lambda > 850$  nm: (i) transfer of vibrational energy from water to oxygen to produce singlet oxygen; and (ii) formation of vibrationally excited solute molecules that react with oxygen.



SINGLET OXYGEN AND RIBOSOMES: INACTIVATION AND  
SITES OF DAMAGE

Harwant Singh, Jeff Bishop and John Merritt

Medical Biophysics Branch  
Atomic Energy of Canada Limited Research Company  
Whiteshell Nuclear Research Establishment  
Pinawa, Manitoba, Canada ROE 1LO

Singlet oxygen has been assigned a significant role in photosensitized processes in biologically important molecules. We have studied the role of singlet oxygen in ribosomes using a number of dyes, psoralens and ketones. With many dyes, evidence has been obtained that singlet oxygen is the inactivation species. However, all of the inactivation seen with some dyes (e.g., acridine orange, methylene blue, etc.) cannot be attributed to singlet oxygen, indicating that other mechanisms are involved. Furthermore, preliminary data with some dyes indicate inactivation in the absence of oxygen (e.g., acridine orange, acridine flavine, etc.). Some dyes (e.g., rhodamine B and ethidium bromide), which reportedly do not cause cell inactivation, cause some inactivation of ribosomes. Inactivation by psoralens is mainly through singlet oxygen. However, singlet oxygen plays a minor role in ketone photosensitized systems, where free radical reactions seem to dominate. Singlet oxygen also appears to be of negligible importance to the inactivation of ribosomes by ionizing radiation.

Analyses of damage to the macromolecules within the ribosomes indicate no chain breaks in ribosomal RNA on photosensitization with the dyes, psoralens or ketones. However, with some dyes, damage to ribosomal proteins is seen by gel electrophoresis.

PREFERENTIAL SOLVATION OF  $O_2^1\Delta_g$  IN MIXED SOLVENTS

Wednesday

B. Stevens<sup>a</sup> and M.A.J. Rodgers<sup>b</sup><sup>a</sup>Dept. of Chemistry, University of South Florida, Tampa, FL 33620<sup>b</sup>CFKR, University of Texas, Austin, TX 78712

A cage displacement model<sup>1</sup>, appropriate to the description of luminescence decay constants ( $1/\tau$ ) at high quencher concentrations where diffusion along concentration gradients is not a quenching prerequisite, is applied to mixed solvent quenching of  $O_2^1\Delta_g$ . The dependence of  $1/\tau$  on mole fractions  $Y_A$  and  $Y_B$  of quenching cosolvents A and B in the solvent cage, is expressed by

$$\frac{1}{\tau} = \frac{1}{\tau_A} + \frac{Y_B \nu_A}{Y_B \nu_A + Y_A \nu_B} \left\{ \frac{1}{\tau_B} - \frac{1}{\tau_A} \right\} \quad I$$

where  $\tau_i$  is the lifetime, and  $\nu_i$  the displacement frequency, of  $O_2^1\Delta_g$  in pure solvent  $i$ .  $Y_A, Y_B$  are related<sup>2</sup> to mole fractions  $X_A, X_B$  in the bulk solvent by

$$Y_A/X_A = (Y_B/X_B) \exp \gamma_A \quad II$$

and insofar as  $\nu_i$  may be expressed in terms of viscosity  $\eta_i$  and molar volume  $V_i$  by<sup>1,3</sup>

$$\nu_i \eta_i V_i = RT \quad III$$

eq I rearranges to

$$f(\tau) = \left( \frac{1}{\tau_B} - \frac{1}{\tau} \right) / \left( \frac{1}{\tau} - \frac{1}{\tau_A} \right) = \frac{X_A \eta_A V_A}{X_B \eta_B V_B} \exp \gamma_A = f(X) \exp \gamma_A \quad IV$$

The index  $\gamma_A$  of preferential solvation of  $O_2^1\Delta_g$  by solvent A in the presence of B, accessible from the experimental quantities  $f(\tau)$  and  $f(X)$ , is found to be essentially zero for a number of polar and non-polar quenching cosolvents; this implies that solvent quenching of  $O_2^1\Delta_g$  is not determined by different stabilities of solvent -  $O_2^1\Delta_g$  complexes.

## Wednesday

1. B. Stevens and K.L. Marsh, J. Phys. Chem., 86 4473 (1982)
2. T. Yoshino, J. Chem. Phys., 24 76 (1956)
3. S. Glasstone, K.J. Laidler and H. Eyring, "The Theory of Rate Processes," McGraw-Hill, New York, 1941.

Wednesday

PRODUCTION AND QUENCHING OF SINGLET OXYGEN BY THE SENSITIZER  
IN THE DYE-SENSITIZED PHOTOOXYGENATIONS

C. TANIELIAN

Laboratoire de Chimie Organique Appliquée,  
Ecole Nationale Supérieure de Chimie, 1, rue Blaise Pascal,  
67008 STRASBOURG - FRANCE

Many dyes are commonly used as singlet oxygen sensitizers (S) in photooxygenation reactions. Nevertheless, there remain some significant gaps in our quantitative knowledge of such fundamental parameters as quantum yield of  $O_2(^1\Delta)$  production  $\gamma_\Delta$  and rate constant  $k_Q$  of quenching of singlet oxygen by S. It is particularly important to realize that there are some situations in which the dye molecules themselves may cause quenching of  $O_2(^1\Delta)$  since neglecting this fact led to erroneous results in the determination of the lifetimes of singlet oxygen and of the rate constant for its reaction with a substrate. This communication reports the measurement of  $\gamma_\Delta$  and  $k_Q$  for various dyes in various solvents saturated with oxygen. For example, it is shown that :

(i) the quantum yield  $\gamma_\Delta$  is 0.50, 0.79 and 0.60 for methylene blue and rose bengal in methanol and chlorophyll a in benzene respectively


(ii) the rate constant of quenching of  $O_2(^1\Delta)$  by rose bengal in acetonitrile is  $k_Q \approx 5 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ .

Harry H. Wasserman


Saturday

Department of Chemistry, Yale University, New Haven, CT 06511

The reactions of singlet oxygen with oxazoles have provided a novel method for protecting carboxyl groups which may then be regenerated in an activated form. Recent use of this procedure in the synthesis of dipeptides will be discussed.



A singlet oxygen reaction which has potential use in synthesis is the oxidation of  $\beta$ -dicarbonyl derivatives leading to vicinal tricarbonyl systems. Thus, while enols of  $\beta$ -diketones show only sluggish reactivity or no reaction with singlet oxygen, the presence of fluoride ion in the reaction medium promotes rapid uptake of singlet oxygen leading to 1,2,3-tricarbonyl products or their hydrates. This reaction most probably results from electron-enrichment in the double bond through hydrogen bonding. Another method for forming vicinal tricarbonyls from  $\beta$ -diketones involves the formation of  $\alpha$ -enamines followed by singlet oxygen oxidative cleavage. Synthetic applications will be discussed.



Laboratory Kinetics and Spectroscopy Applied to Airglow Studies


R. P. Wayne

Physical Chemistry Laboratory, South Parks Road, Oxford, OX1 3QZ, U.K.


This survey of excitation processes for singlet oxygen in planetary atmospheres is intended to provide an introduction to the following papers which discuss in more detail laboratory investigations and field measurements. The three singlet states,  $a^1\Delta_g$ ,  $b^1\Sigma_g^+$  and  $c^1\Sigma_u^-$  contribute to features in the airglows of Venus, Earth and Mars, the most intense feature being in each case the dayglow of the  $O_2(a^1\Delta_g - x^3\Sigma_g^-)$  'Infrared Atmospheric Band'. Ozone photolysis is the daytime source of  $O_2(a^1\Delta_g)$  on Earth and Mars, but is unlikely to be on Venus. Some remaining problems about the ozone photolytic source are reviewed, and possible excitation mechanisms on Venus are considered.  $O_2(b^1\Sigma_g^+)$  is excited during the day in the Terrestrial atmosphere by a combination of resonance scattering and of energy transfer from  $O(^1D)$  to  $O_2$ . The state is unimportant on either Venus or Mars because of its sensitivity to quenching by  $CO_2$ , the major atmospheric constituent.

Nightglow emissions from  $O_2(a^1\Delta_g)$  and  $O_2(b^1\Sigma_g^+)$  are much harder to explain than the daytime phenomena, because they must be excited from chemical energy stored in species, such as H, O or  $O_3$ , that persist at night. Of the many mechanisms that have been proposed, none has been substantiated explicitly by laboratory experiments, and the hypothetical sources are frequently judged by their ability to match atmospheric measurements of airglow both for absolute intensity and for altitude profile. We consider here the additional factor of the plausibility of the mechanisms.

A particularly important question is whether recombination of ground state atomic oxygen



populates the several molecular states, with which the atoms correlate, statistically, or whether some states (perhaps even the ground state) are formed preferentially. Several developments, of which we shall hear more in later lectures, have focussed interest on this problem. For a start, the discovery that  $O_2(a^1\Delta_g)$  is formed by the recombination of O on pyrex walls has indirectly made it practically certain that the atomic oxygen 'green line' in the airglow is excited by a two-step 'Barth' mechanism. Thus some state  $O_2^\#$  excited in reaction (1) itself reacts with O



Such mechanisms can, at least on paper, explain the intensities, peak emission altitudes, and layer thicknesses of  $O(^1S)$ ,  $O_2(a^1\Delta_g)$  and  $O_2(b^1\Sigma_g^+)$  emissions,

although the states #, † and †† of  $O_2$  have not yet been identified. Experimental confirmation of a reaction corresponding to process (4) will be presented later in this meeting.

In the context of these ideas, recent advances in our understanding of the  $c\ ^1\Sigma_u^-$  state of  $O_2$ , both in atmospheres and in the laboratory, may be relevant. Following the identification of the 'Herzberg II'  $O_2(c\ ^1\Sigma_u^- - X\ ^3\Sigma_g^-)$  system in the Venusian airglow, laboratory investigations showed that the  $c\ ^1\Sigma_u^-$  state is formed in the O atom recombination. At first, it appeared that  $CO_2$  was a necessary third body M for the appearance of the state, but it then emerged that  $O_2(c\ ^1\Sigma_u^-)$  was formed, but in reduced yield, with other third bodies. Clever kinetic studies, in skilled hands, have indicated that O atoms themselves quench  $O_2(c\ ^1\Sigma_u^-)$ , and also that the peculiar influence of  $CO_2$  may possibly arise from its ability to funnel most of the recombining pairs of atoms to the  $c\ ^1\Sigma_u^-$  state. The existence of the laboratory source of  $O_2(c\ ^1\Sigma_u^-)$  has made it possible to obtain high resolution spectra of the Herzberg II system. In turn, comparison of these spectra with those of the Earth's airglow have revealed that  $O_2\ c\ ^1\Sigma_u^-$  is indeed present in our atmosphere. Two differences distinguish Earth's Herzberg II emission from that of Venus : it is about 27 times weaker, but it does exhibit transitions from vibrationally excited levels, with the peak emission corresponding to  $v' = 7$ . The relative paucity of  $CO_2$  in the Earth's atmosphere can explain both observations, if  $CO_2$  is efficient in promoting population of the electronic state as well as in vibrational quenching. More subtle arguments further show that the relative intensities of the Herzberg II system on Venus and Earth, and the virtual absence on Mars, can be related to the concentrations of  $CO_2$ ,  $N_2$  or  $O_2$  compared to that of O in the atmospheres of the three planets.

Finally, we are forced to ask whether the states of  $O_2$  designated as #, †, †† actually correspond to  $O_2(c\ ^1\Sigma_u^-)$  possessing differing degrees of vibrational excitation. Energy requirements are satisfied for reaction (2) for  $O_2(c\ ^1\Sigma_u^-)$  with  $v'$  greater than ~2 or 3. Reactions (3) and (4) are thermally accessible for  $v' = 0$ . Alternative states for  $O_2^\#$ , such as the  $A\ ^2\Sigma_u^-$  or  $A'\ ^3\Delta_u$ , are made less likely by the weakness of  $O(^1S)$  emission in the Venusian airglow. The  $v' = 0$  levels of c, A, and A' states are all populated in the Venusian atmosphere, but only that of the c state lies below the  $O(^1S)$  excitation energy and thus has an activation energy that would make the process slow at the altitude of peak emission. On the other hand, the new evidence about reaction (4) suggests that, even though an energy exchange process excite  $O_2(b\ ^1\Sigma_g^+)$ , none of the states  $c\ ^1\Sigma_u^-$ ,  $A\ ^3\Sigma_u^-$  or  $A'\ ^3\Delta_u$  can be involved, at least in their lower vibrational levels.

Friday

IF  $B^3\Pi(0^+)$  Pumping by Energy Transfer from  $O_2^1\Delta_g$  to IFX $^1\Sigma^+$ 




## A Potential Chemical Laser Pumping Scheme.\*

P. D. Whitefield

McDonnell Douglas Research Laboratories

St. Louis, MO 63166

Studies have been made of the IF (B-X) chemiluminescence resulting from the mixing of three sets of reagents:



Simultaneous mixing of the third set of reagents produced the most intense IF (B-X) chemiluminescence. In all cases the chemiluminescence indicated that rapid rotational thermalization occurs. When  $O_2^1\Delta_g$  was present, however, an extremely non-thermal vibrational population in the IF  $B^3\Pi(0^+)$  was observed.

Indirect evidence suggests that sequential pumping of ground state IF(X) molecules produces the observed chemiluminescence. A possible reaction scheme which describes a sequential pumping mechanism will be presented.

The photon yield measured for the chemiluminescence arising from the third set of reagents is small,  $\sim 0.3\%$ . This yield can be explained largely by the low densities of  $O_2^1\Delta_g$  used in these studies. The  $I_2 + F + O_2^1\Delta_g$

\*This research is supported by AFOSR and Independent Research and Development funds from McDonnell Douglas Research Laboratories.



system may hold promise as an efficient chemical excitation source for IF (B), particularly if higher densities of  $O_2^1\Delta_g$  can be produced and utilized.

Finally, collisional ladder climbing has been observed when the B states of the analogous interhalogens BrF and BrCl have been excited.<sup>1,2</sup> Ladder climbing from vibrational levels close to the predissociation limit result in non-radiative loss to predissociation inhibiting the establishment of a population inversion. No measurements have been published to date on ladder climbing in the IF(B) state, but inferences of its existence from the chemiluminescence data will be discussed.

#### References

1. M. A. A. Clyne and I. S. McDermid, J. Chem. Soc. Faraday Trans. 274, 1376, (1978).
2. M. A. A. Clyne and I. S. McDermid, Faraday Discuss. 67, 316, (1979).

SELF-SENSITIZED PHOTOOXIDATION OF PROTOPORPHYRIN IX AND RELATED  
PORPHYRINS IN ERYTHROCYTE GHOSTS AND MICROEMULSIONS: A NOVEL  
PHOTOOXIDATION PATHWAY

Marianne Krieg and David G. Whitten

Departments of Chemistry

University of North Carolina, Chapel Hill, NC 27514

University of Rochester, Rochester, New York 14627

**ABSTRACT:** Previous studies have shown that protoporphyrin IX can sensitize its own photooxidation by paths involving primarily singlet oxygen and to a lesser extent superoxide (via excited state electron transfer). These reactions involve reaction of the vinyl groups of the protoporphyrin to yield porphyrins with modified side chains. Quantum efficiencies are generally low and the product distribution is somewhat solvent dependent. In recent work we have examined the photooxidation of protoporphyrin IX, mesoporphyrin IX and hematoporphyrin IX in erythrocyte "ghosts", a natural membrane system containing saturated lipids, unsaturated lipids and a full complement of membrane proteins. We observe that all three porphyrins are rapidly photooxidized in the "ghosts" and that the products produced from protoporphyrin IX do not include the "usual" singlet oxygen products. We have been able to model the kinetic behavior observed in the natural membrane systems by using an oil/water microemulsion as a solvent medium and adding various amino acids that are easily oxidized. In particular we have investigated the behavior of methionine and a number of related thioether derivatives. The results of this study suggest that the porphyrins sensitize singlet oxygen efficiently but that the

singlet oxygen is rapidly scavenged by substrates such as methionine and other amino acids. The oxygenated amino acids can subsequently act as agents to efficiently oxidize the porphyrins by attacking the porphyrin ring directly. Thus, although singlet oxygen is clearly indicated to be involved in these reactions, the actual agent in the photooxidation of the porphyrins appears to be an intermediate occurring subsequent to its consumption.

THE MECHANISM OF QUENCHING OF  
SINGLET OXYGEN BY MOLECULAR IODINE

by

F. Wilkinson and A. Farmilo

Department of Chemistry  
Loughborough University of Technology  
LOUGHBOROUGH  
Leicestershire. LE11 3TU. England

The various methods for determining singlet oxygen quenching constants are reviewed in Reference 1 which includes a compilation of quenching constants by 690 compounds. Although quenching of singlet oxygen by some alkyl iodides and the iodide ion has been studied<sup>(1)</sup> quenching by molecular iodine has not been reported previously despite the fact that  $I_2$  quenches excited singlet<sup>(2)</sup> and excited triplet states<sup>(3)</sup> of aromatic compounds with high efficiency. We have used a nanosecond pulsed ruby laser to sensitize the production of singlet oxygen using anthracene and other organic triplet states as singlet oxygen sensitizers and determined the pseudo first-order decay constant of singlet oxygen  $k_D$  from a kinetic analysis of the disappearance of a small proportion of a fixed concentration of the singlet oxygen acceptor diphenylisobenzofuran, DPBF. In the presence of various amounts of quencher

$$k_D = k_1 + k_R[DPBF]_{AV} + k_Q[Q]$$

where  $k_1$  is the decay constant for singlet oxygen in the absence of DPBF and any quencher  $Q$  and  $k_R$  and  $k_Q$  are the second-order reaction and quenching constants. Thus a plot of  $k_D$  versus  $[Q]$  is often linear and  $k_Q$  can be obtained from the slope (see refs. 1 and 4). In the case of iodine as a potential quencher the plot of  $k_D$  versus  $[Q]$  is not linear although the initial slope gives a  $k_Q$  value of  $1.4 \times 10^9 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ . The mechanism for quenching and the reasons for the non-linear plot of  $k_D$  versus  $[I_2]$  will be discussed.

REFERENCES

1. F. Wilkinson and J.G. Brummer, J.Phys.and Chem.Ref.Data, 1981, 10, 809.
2. F. Costano, A. Lozaro, M. Martinez and E. Martinez, Proc.of IX IUPAC Symposium de Photochemie, Pau, 1982, p.58.
3. G. Porter and F. Wilkinson, Proc.Roy.Soc., 1961, 264A, 1.
4. F. Wilkinson and A. Farmilo, Photochem.Photobiol., 1983, 18, 447.

Thursday

ACTIVE OXYGEN CHEMISTRY OF OCULAR TISSUES- S. Zigman  
Ophthalmology Research Laboratory, University of Rochester  
Rochester, N. Y. 14642

Because they must receive and absorb much radiant energy to perform their visual functions, ocular tissues must also be capable of protecting themselves against free radicals and oxidation products that would destroy them. While only low levels and limited wavelengths of light are required for vision, excessive quantities and visually useless light enter the eye and is absorbed by various tissues whose function is to deliver the high quanta necessary for vision to occur. Besides these considerations, it is important to note that the levels of oxygen vary tremendously in these tissues, with very low levels being present in the ocular lens and very high levels in the retina.

Data on the spectral quality and quantities of radiant energy interacting with various ocular tissues, their oxygen content, and enzymes and anti-oxidants present to scavenge or protect against photooxidative damage needs to be collated in order to assess the potential hazard to these tissues relative to oxygen excited states and radicals. Such data will be provided in our presentation. Besides this information chemopathological conditions have been observed in the ocular tissues (ie: cataract; retinal degeneration) that have been ascribed to imbalances in the above-mentioned system. These will be described and the potential that oxygen excited states and free radicals have an important role in the damages stated will be discussed.

This talk will introduce practical situations relating eye health to the handling of oxygen products induced by radiant energy and metabolism.

# AUTHOR INDEX

Acs, A.	B3	Fink, E.H.	<u>F1</u>
Agarwal, S.K.	M3	Foot, C.S.	<u>F2</u> , P3
Ando, W.	<u>A1</u>	Forbes, P.D.	L2
Antonsen, S.A.	S5	Forsskah1, I.	<u>F3</u>
Arepalli, S.	H2	Frimer, A.A.	<u>F4</u>
Bishop, J.	S6	Gabriel, R.	B3
Bogan, D.J.	<u>B1</u>	Gandhi, P.B.	<u>G1</u>
Boodaghians, R.	B2	Ghaeni, S.	B3
Borrell, P.	<u>B2</u>	Gharapetian, H.M.	G5
Borrell, P.M.	B2	Gorman, A.A.	<u>G2</u>
Brauer, H.-D.	<u>B3</u> , S4	Griffin, G.W.	<u>G3</u>
Brunet, J.E.	D1	Grossweiner, L.I.	<u>G4</u>
Butler, R.	E1	Guillet, J.E.	<u>G5</u>
Campa, A.	D1	Gurinovich, G.P.	<u>G6</u>
Chowdhary, R.K.	C2	Hall, G.E.	H2
Connolly, J.S.	<u>C1</u>	Hamblett, I.	G2
Cundall, R.B.	<u>C2</u>	Heidner, R.F. III	<u>H1</u>
Davies, A.K.	C2	Houston, P.L.	<u>H2</u>
Davies, R.E.	L2	Inoue, K.	S1
Dobrowolski, D.C.	F2	Jefford, C.W.	<u>J1</u>
Dobrezelewski, M.	M2	Kabe, Y.	A1
Drews, W.	B3	Kanofsky, J.R.	<u>K1</u>
Dubey, R.D.	G1	Kessick, R.F.	M2
Duran, N.	<u>D1</u>	Khan, A.U.	<u>K2</u>
Dzhagarov, B.M.	<u>D2</u>	Kirschenheuter, G.P.	G3
Eisenberg, W.C.	<u>E1</u>	Koroll, G.W.	S5
Evans, D.F.	<u>E2</u>	Krasnovsky, A.A.	<u>K3</u>
Faler, G.R.	S3	Krieg, M.	W4
Faljori-Alario, A.	D1	Kruse, H.	F1
Farias-Furtado, S.T.	D1	Llewellyn, E.J.	<u>L1</u>
Farmilo, A.	W5	Logani, M.K.	<u>L2</u>
		Lopez, L.	S2

## AUTHOR INDEX

-2-

MacInnis, K.	G5	Siddiqui, S.	S2
Maier, M.	<u>M1</u>	Singh, A.	<u>S5</u>
Marinelli, W.	H2	Singh, H.	<u>S6</u>
Marsh, K.L.	C1	Snelson, A.	E1
Matsuura, T.	S1	Stanbro, W.D.	P1
McDade, I.C.	L1	Stevens, B.	<u>S7</u>
Mendenhall, G.D.	<u>M2</u>	Sugiyama, H.	S1
Merritt, J.	S6		
Murray, R.W.	E1, <u>M3</u>	Takata, T.	A1
		Tanielian, C.	<u>T1</u>
Nultsch, W.	<u>N1</u>	Taylor, K.	E1
		Trost, W.	B3
Ogryzlo, E.A.	<u>O1</u>		
		Ueno, K.	A1
Parker, J.G.	<u>P1</u> , <u>P2</u>	Umrigar, P.P.	G3
Peak, J.G.	P3	Upton, M.W.	E2
Peak, M.J.	<u>P3</u>		
Prasad, G.	S2	Villasenor, S.R.	S3
Rabek, J.F.	<u>R1</u>	Wassell, P.T.	O1
Ranby, B.	R1	Wasserman, H.H.	<u>W1</u>
Richards, D.S.	B2	Wayne, R.P.	<u>W2</u>
Rodgers, M.A.J.	G2, <u>R2</u> , S7	Whitefield, P.D.	<u>W3</u>
Russell, V.P.	C2	Whitten, D.G.	<u>W4</u>
		Wiesenfeld, J.R.	H2
Saito, I.	<u>S1</u>	Wildt, J.	F1
Salokhiddinov, K.I.	D2	Wilkinson, F.	<u>W5</u>
Schaap, A.P.	<u>S2</u> , <u>S3</u>	Winter, R.	F1
Schaffner, K.	B3		
Schmidt, R.	B3, <u>S4</u>	Young, A.T.	H2
Schuchart, H.	<u>N1</u>		
Shen, Y.Q.	O1	Zabel, F.	F1
Sherren, J.	G5	Zigman, S.	<u>Z1</u>

**END**

**FILMED**

**12-84**

**DTIC**